

t1\_relset\_1 (TMRevaS-  
BkgGM6rTVu1FXQj9PMc7F9gdprEJ)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1\_tarski X0 X1) \wedge (r1\_tarski X1 X2)) \Rightarrow (r1\_tarski X0 X2) \quad (2)$$

**Theorem 1**

$$\forall X0. \forall X1. \forall X2. \forall X3. (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X1 X2))) \Rightarrow ((r1\_tarski X0 X3) \Rightarrow (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X1 X2))))$$