

t1\_rewrite3  
(TMFKE2gjSbcf7rv96eVQsRsCMCbXmxzo6nB)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k4\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k1\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (( \\ v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow (\forall X2.(X0 \in k1\_relset\_1 \\ k5\_numbers X1) \Rightarrow (k1\_funct\_1 (k7\_finseq\_1 (k9\_finseq\_1 X2) X1) \\ (k1\_nat\_1 X0 np\_1) = k1\_funct\_1 X1 X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k4\_finseq\_1 X0 = k9\_xtuple\_0 X0) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0) \wedge (m1\_subset\_1 X1 k5\_numbers)) \Rightarrow (k1\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & ((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 \ k5\_numbers) \wedge ((v1\_funct\_1 X0) \wedge \\ & (v1\_finseq\_1 X0)))) \end{aligned} \quad (6)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. \forall X1. (v7\_ordinal1 X1) \Rightarrow (\forall X2. ((v1\_relat\_1 \\ & X2) \wedge ((v1\_funct\_1 X2) \wedge (v1\_finseq\_1 X2))) \Rightarrow ((X1 \in k4\_finseq\_1 X2) \Rightarrow \\ & (k1\_funct\_1 (k7\_finseq\_1 (k9\_finseq\_1 X0) X2) (k2\_xcmplx\_0 X1 \\ & np\_1) = k1\_funct\_1 X2 X1))) \end{aligned}$$