

t1_simplex0 (TMZ- eRQKNT42MnSbrxaEcDNDUvzQLm5Vhu4Q)

October 27, 2020

Let $v1_classes1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_setfam_1 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. k9_setfam_1 X0 = k1_zfmisc_1 X0 \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_zfmisc_1 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (r1_tarski X2 X0)) \quad (3)$$

Assume the following.

$$\forall X0. (v1_classes1 X0) \Leftrightarrow (\forall X1. \forall X2. ((X1 \in X0) \wedge (r1_tarski X2 X1)) \Rightarrow (X2 \in X0)) \quad (4)$$

Theorem 1

$$\forall X0. (v1_classes1 X0) \Leftrightarrow (\forall X1. (X1 \in X0) \Rightarrow (r1_tarski (k9_setfam_1 X1) X0))$$