

t1_t_0topsp
(TMabCX2wd8dfpvYUqrrqJkgjQCQUjjNKmyh6)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k8_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \neg(v1_xboole_0 X0) \wedge ((X0 \neq X1) \wedge (v1_xboole_0 X1)) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1_funct_1 X3) \wedge \\ & ((v1_funct_2 X3 X0 X1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))) \Rightarrow (\forall X4. \neg(X4 \in k7_relset_1 X0 X1 X3 X2) \wedge (\forall X5. \\ & \neg(X5 \in X0) \wedge ((X5 \in X2) \wedge (X4 = k1_funct_1 X3 X5)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1_funct_1 X3) \wedge \\ & ((v1_funct_2 X3 X0 X1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))) \Rightarrow ((r1_tarski X2 X0) \Rightarrow (((X1 = k1_xboole_0) \wedge (X0 \neq k1_xboole_0)) \vee \\ & (r1_tarski X2 (k8_relset_1 X0 X1 X3 (k7_relset_1 X0 X1 X3 X2)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(k8_relset_1 X0 X1 X2 X3 = k8_relat_1 X2 X3) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(k7_relset_1 X0 X1 X2 X3 = k7_relat_1 X2 X3) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0)\wedge(((v1_funct_1 X2)\wedge((v1_funct_2 X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\wedge(m1_subset_1 X3 X0))))\Rightarrow(k3_funct_2 X0 X1 X2 X3 = k1_funct_1 X2 X3) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\exists X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\wedge((v1_xboole_0 X2)\wedge((v1_relat_1 X2)\wedge((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)))) \quad (9)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(m1_subset_1 (k8_relset_1 X0 X1 X2 X3) (k1_zfmisc_1 X0)) \quad (11)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.\forall X2.(X2 = k8_relat_1 X0 X1)\Leftrightarrow(\forall X3.(X3 \in X2)\Leftrightarrow((X3 \in k9_xtuple_0 X0)\wedge(k1_funct_1 X0 X3 \in X1)))) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow(X2 \in X1)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(X0 = X1)\Leftrightarrow((r1_tarski X0 X1)\wedge(r1_tarski X1 X0)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (15)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow \\ & (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow (\forall X3.(m1_subset_1 \\ & X3 (k1_zfmisc_1 X0)) \Rightarrow ((\forall X4.(m1_subset_1 X4 X0) \Rightarrow (\forall X5. \\ & (m1_subset_1 X5 X0) \Rightarrow (((X4 \in X3) \wedge (k3_funct_2 X0 X1 X2 X4 = k3_funct_2 \\ & X0 X1 X2 X5)) \Rightarrow (X5 \in X3)))) \Rightarrow (k8_relset_1 X0 X1 X2 (k7_relset_1 X0 X1 \\ & X2 X3) = X3)))))) \end{aligned}$$