

t1_tex_2

(TMM5nFvacxyNfQpepGaF35k9Mn84KcLe7cz)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_card_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_zfmisc_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 (k1_tarski X1)) \Leftrightarrow ((X0 = k1_xboole_0) \vee (X0 = k1_tarski X1)) \quad (1)$$

Assume the following.

$$\forall X0. \neg(\neg v1_xboole_0 X0) \wedge ((v1_zfmisc_1 X0) \wedge (\forall X1. X0 \neq k1_tarski X1)) \quad (2)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (3)$$

Assume the following.

$$\forall X0. (v3_card_1 X0 np_1) \Rightarrow ((\neg v1_xboole_0 X0) \wedge (v1_zfmisc_1 X0)) \quad (4)$$

Theorem 1

$$\forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (v3_card_1 X1 np_1) \Rightarrow ((r1_tarski X0 X1) \Rightarrow (X0 = X1)))$$