

t1_topgen_3
(TMbEZsJvbseRUPxGdtQd9yab16q4BYVPEag)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v3_abian : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_setfam_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow ((v3_abian X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Leftrightarrow (k5_setfam_1 X0 X1 = X0)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow (k5_setfam_1 X0 X1 = k3_tarski X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\forall X2. (X2 \in X1) \Rightarrow (X2 \in X0)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k3_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. (X2 \in X3) \wedge (X3 \in X0))) \quad (5)$$

Theorem 1

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow ((v3_abian X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Leftrightarrow (\forall X2. \neg (X2 \in X0) \wedge (\forall X3. (m1_subset_1 X3 (k1_zfmisc_1 X0)) \Rightarrow (\neg (X3 \in X1) \wedge (X2 \in X3))))))$$