

t1_topreal7

(TMYi3GnxVt81XYmD6juTvL2unY9Ex6CY5Bp)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_square_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k4_xxreal_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v2_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & X0 X1) \wedge (r1_xxreal_0 X2 X3)) \Rightarrow (r1_xxreal_0 (k2_xcmplx_0 X0 X2) (\\ & k2_xcmplx_0 X1 X3)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X2 X1)) \Rightarrow \\ & (r1_xxreal_0 (k4_xxreal_0 X0 X2) X1)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (r1_xxreal_0 X0 (k4_xxreal_0 X0 X1))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (k7_real_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (m1_subset_1 X1 k1_numbers)) \Rightarrow (k2_square_1 X0 X1 = k4_xxreal_0 X0 X1) \quad (5)$$

Assume the following.

$$v3_membered k1_numbers \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers)\wedge(v1_xreal_0 X1))\Rightarrow(m1_subset_1 (k7_real_1 X0 X1) k1_numbers) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers)\wedge(m1_subset_1 X1 k1_numbers))\Rightarrow(m1_subset_1 (k2_square_1 X0 X1) k1_numbers) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0)\wedge(v1_xxreal_0 X1))\Rightarrow(k4_xxreal_0 X0 X1 = k4_xxreal_0 X1 X0) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xxreal_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.(v3_membered X0)\Rightarrow(v2_membered X0) \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (12)$$

Assume the following.

$$\forall X0.(v2_membered X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(v1_xxreal_0 X1)) \quad (13)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(\forall X1.(m1_subset_1 X1 k1_numbers)\Rightarrow(\forall X2.(m1_subset_1 X2 k1_numbers)\Rightarrow(\forall X3.(m1_subset_1 X3 k1_numbers)\Rightarrow(r1_xxreal_0 (k2_square_1 (k7_real_1 X0 X2) (k7_real_1 X1 X3)) (k7_real_1 (k2_square_1 X0 X1) (k2_square_1 X2 X3))))))$$