

t1\_toprns\_1 (TMSeXHGR-  
bGV8fVz6nBx5c9yhoxYTMmPm2Qs)

October 27, 2020

Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k15\_euclid : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $l1\_struct\_0 : \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v5\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $l1\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $l1\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $l1\_pre\_topc : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 X1) \Rightarrow ((v1\_xboole\_0 X1) \vee (X0 \in X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1\_subset\_1 X0 X1) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. (( \\ \neg v2\_struct\_0 X1) \wedge (l1\_struct\_0 X1)) \Rightarrow (\forall X2. (m1\_subset\_1 \\ X2 (u1\_struct\_0 X1)) \Rightarrow (((v1\_funct\_1 X0) \wedge ((v1\_funct\_2 X0 k5\_numbers \\ (u1\_struct\_0 X1)) \wedge (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ k5\_numbers (u1\_struct\_0 X1)))))) \Leftrightarrow ((k9\_xtuple\_0 X0 = k5\_numbers) \wedge \\ (\forall X3. (X3 \in k5\_numbers) \Rightarrow (m1\_subset\_1 (k1\_funct\_1 X0 X3) \\ (u1\_struct\_0 X1))))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2. (m2\_subset\_1 \\ X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \quad (4)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (5)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1) \wedge (v3\_ordinal1 k4\_ordinal1) \quad (6)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow ((\neg v2\_struct\_0 (k15\_euclid X0)) \wedge (v5\_rltopsp1 (k15\_euclid X0))) \quad (7)$$

Assume the following.

$$\neg v1\_xboole\_0 k1\_numbers \quad (8)$$

Assume the following.

$$\forall X0.\exists X1.m1\_subset\_1 X1 X0 \quad (9)$$

Assume the following.

$$\forall X0.(l1\_rltopsp1 X0) \Rightarrow ((l1\_rlvect\_1 X0) \wedge (l1\_pre\_topc X0)) \quad (10)$$

Assume the following.

$$\forall X0.(l1\_pre\_topc X0) \Rightarrow (l1\_struct\_0 X0) \quad (11)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (12)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow ((v5\_rltopsp1 (k15\_euclid X0)) \wedge (l1\_rltopsp1 (k15\_euclid X0))) \quad (13)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (14)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(m2\_subset\_1 X0 k1\_numbers k5\_numbers) \Rightarrow (\forall X1. \\ & ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (((v1\_funct\_1 X1) \wedge (v1\_funct\_2 \\ & X1 k5\_numbers (u1\_struct\_0 (k15\_euclid X0))) \wedge (m1\_subset\_1 X1 \\ & (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers (u1\_struct\_0 (k15\_euclid \\ & X0)))))) \Leftrightarrow ((k9\_xtuple\_0 X1 = k5\_numbers) \wedge (\forall X2.(m2\_subset\_1 \\ & X2 k1\_numbers k5\_numbers) \Rightarrow (m1\_subset\_1 (k1\_funct\_1 X1 X2) (u1\_struct\_0 \\ & (k15\_euclid X0)))))) \end{aligned}$$