

t20_arytm_0
(TMJdyKi5uG7TmnXpNTZHtMwGJwhYDsFyBBs)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k2_arytm_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_arytm_0 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow ((X1 = np_1) \Rightarrow (k2_arytm_0 X0 X1 = X0))) \quad (1)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow (\forall X2.(m1_subset_1 X2 k1_numbers) \Rightarrow (k2_arytm_0 X0 (k2_arytm_0 X1 X2) = k2_arytm_0 (k2_arytm_0 X0 X1) X2))) \quad (2)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (m1_subset_1 (k4_arytm_0 X0) k1_numbers) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (m1_subset_1 X1 k1_numbers)) \Rightarrow (m1_subset_1 (k2_arytm_0 X0 X1) k1_numbers) \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow (((X0 \neq k6_numbers) \Rightarrow ((X1 = k4_arytm_0 X0) \Leftrightarrow (k2_arytm_0 X0 X1 = np_1))) \wedge ((X0 = k6_numbers) \Rightarrow ((X1 = k4_arytm_0 X0) \Leftrightarrow (X1 = k6_numbers))))) \quad (5)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow ((X1 \neq k6_numbers) \Rightarrow (k2_arytm_0 (k2_arytm_0 X0 X1) (k4_arytm_0 X1) = X0)))$$