

## t20\_bvfunc\_6

(TMZXJ5vuFUMZU86z9mU3SFCbnLacPrz69Qg)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_margrel1 : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r2\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k12\_bvfunc\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge ( \\ & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge \\ & ((v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (r2\_funct\_2 X0 k6\_margrel1 \\ & (k9\_bvfunc\_1 X0 X1 X2) (k5\_bvfunc\_1 X0 (k1\_bvfunc\_1 X0 X1) X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge ( \\ & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow ((r2\_funct\_2 X0 k6\_margrel1 \\ & (k9\_bvfunc\_1 X0 (k12\_bvfunc\_1 X0) X1) (k12\_bvfunc\_1 X0)) \Rightarrow (r2\_funct\_2 \\ & X0 k6\_margrel1 X1 (k12\_bvfunc\_1 X0)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.(((v1\_funct\_1 X2) \wedge \\ & ((v1\_funct\_2 X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 X1)))))) \wedge ((v1\_funct\_1 X3) \wedge ((v1\_funct\_2 X3 X0 X1) \wedge (m1\_subset\_1 \\ & X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))))) \Rightarrow ((r2\_funct\_2 X0 X1 X2 \\ & X3) \Leftrightarrow (X2 = X3)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge((v1\_funct\_1 X1)\wedge((v1\_funct\_2 X1 X0 k6\_margrel1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1))))))\Rightarrow(k1\_bvfunc\_1 X0 (k1\_bvfunc\_1 X0 X1) = X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0)\wedge(((v1\_funct\_1 X1)\wedge((v1\_funct\_2 X1 X0 k6\_margrel1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1))))))\wedge((v1\_funct\_1 X2)\wedge((v1\_funct\_2 X2 X0 k6\_margrel1)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))))\Rightarrow(k5\_bvfunc\_1 X0 X1 X1 = X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0)\wedge(((v1\_funct\_1 X1)\wedge((v1\_funct\_2 X1 X0 k6\_margrel1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1))))))\wedge((v1\_funct\_1 X2)\wedge((v1\_funct\_2 X2 X0 k6\_margrel1)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))))\Rightarrow((v1\_funct\_1 (k5\_bvfunc\_1 X0 X1 X2))\wedge((v1\_funct\_2 (k5\_bvfunc\_1 X0 X1 X2) X0 k6\_margrel1)\wedge(m1\_subset\_1 (k5\_bvfunc\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge((v1\_funct\_1 X1)\wedge((v1\_funct\_2 X1 X0 k6\_margrel1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1))))))\Rightarrow((v1\_funct\_1 (k1\_bvfunc\_1 X0 X1))\wedge((v1\_funct\_2 (k1\_bvfunc\_1 X0 X1) X0 k6\_margrel1)\wedge(m1\_subset\_1 (k1\_bvfunc\_1 X0 X1) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \quad (7)$$

Assume the following.

$$\forall X0.(\neg v1\_xboole\_0 X0)\Rightarrow((v1\_funct\_1 (k12\_bvfunc\_1 X0))\wedge((v1\_funct\_2 (k12\_bvfunc\_1 X0) X0 k6\_margrel1)\wedge(m1\_subset\_1 (k12\_bvfunc\_1 X0) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \quad (8)$$

**Theorem 1**

$$\forall X0.(\neg v1\_xboole\_0 X0)\Rightarrow(\forall X1.((v1\_funct\_1 X1)\wedge((v1\_funct\_2 X1 X0 k6\_margrel1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1))))))\Rightarrow(\forall X2.((v1\_funct\_1 X2)\wedge((v1\_funct\_2 X2 X0 k6\_margrel1)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1))))))\Rightarrow(((r2\_funct\_2 X0 k6\_margrel1 (k5\_bvfunc\_1 X0 X1 X2) (k12\_bvfunc\_1 X0))\wedge(r2\_funct\_2 X0 k6\_margrel1 (k1\_bvfunc\_1 X0 X1) (k12\_bvfunc\_1 X0)))\Rightarrow(r2\_funct\_2 X0 k6\_margrel1 X2 (k12\_bvfunc\_1 X0))))$$