

t20_cqc_the1
(TMQikc94Y24Y5zfPgizM5UtDYCAgE7zfHit)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $v1_cqc_the1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_cqc_the1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (k3_cqc_lang X0))) \Rightarrow (r1_tarski X1 (k1_cqc_the1 X0 X1))) \quad (1)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (k3_cqc_lang X0))) \Rightarrow (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k3_cqc_lang X0))) \Rightarrow ((v1_cqc_the1 X1 X0) \wedge (r1_tarski X2 X1)) \Rightarrow (r1_tarski (k1_cqc_the1 X0 X2) X1)))) \quad (2)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (k3_cqc_lang X0))) \Rightarrow (v1_cqc_the1 (k1_cqc_the1 X0 X1) X0)) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(X0 = X1) \Leftrightarrow ((r1_tarski X0 X1) \wedge (r1_tarski X1 X0)) \quad (4)$$

Theorem 1

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (k3_cqc_lang X0))) \Rightarrow ((v1_cqc_the1 X1 X0) \Leftrightarrow (k1_cqc_the1 X0 X1 = X1)))$$