

t20_euclid_4

(TMZRdhcPv4tevHbmNo4fhKj2q8wfyYDApUbq)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $k23_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_euclid : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v3_finseq_1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v3_card_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_funct_1 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 \\ & X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 \\ & X1) \wedge ((v3_valued_0 X1) \wedge (v1_finseq_1 X1)))) \Rightarrow (\forall X2.((v1_relat_1 \\ & X2) \wedge ((v1_funct_1 X2) \wedge ((v3_valued_0 X2) \wedge (v1_finseq_1 X2)))) \Rightarrow \\ & (((k3_finseq_1 X0 = k3_finseq_1 X1) \wedge (k3_finseq_1 X1 = k3_finseq_1 \\ & X2)) \Rightarrow (k23_rvsum_1 (k4_rvsum_1 X0 X1) X2 = k9_binop_2 (k23_rvsum_1 \\ & X0 X2) (k23_rvsum_1 X1 X2)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_2 X1 X0) \Rightarrow (\forall X2. (m2_finseq_2 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (k9_binop_2 X0 X1 = k2_xcmplx_0 X0 X1) \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers)\wedge(v1_xreal_0 X1))\Rightarrow(k7_real_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7_ordinal1 X0)\wedge((m1_subset_1 X1 (k1_euclid X0))\wedge(m1_subset_1 X2 (k1_euclid X0))))\Rightarrow(k7_euclid X0 X1 X2 = k1_valued_1 X1 X2) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v3_valued_0 X0)\wedge(v1_finseq_1 X0))))\wedge((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge((v3_valued_0 X1)\wedge(v1_finseq_1 X1))))))\Rightarrow(k4_rvsum_1 X0 X1 = k1_valued_1 X0 X1) \quad (7)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow(k3_finseq_1 X0 = k1_card_1 X0) \quad (8)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v3_finseq_1 (k1_euclid X0)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2 X1 X0)\Rightarrow(\forall X2.(m2_finseq_2 X2 X0 X1)\Rightarrow(m2_finseq_1 X2 X0)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v3_valued_0 X0)\wedge(v1_finseq_1 X0))))\wedge((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge((v3_valued_0 X1)\wedge(v1_finseq_1 X1))))))\Rightarrow(m1_subset_1 (k23_rvsum_1 X0 X1) k1_numbers) \quad (12)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(m1_finseq_2 (k1_euclid X0) k1_numbers) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(v3_card_1 X1 X0)\Leftrightarrow(k1_card_1 X1 = X0) \quad (14)$$

Assume the following.

$$\forall X0.(v3_finseq_1 X0)\Rightarrow(v4_funct_1 X0) \quad (15)$$

Assume the following.

$$\forall X0.(v4_funct_1 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(v1_relat_1 X1)\wedge(v1_funct_1 X1)) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow(v5_relat_1 X1 X0) \quad (17)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v5_relat_1 X0 k1_numbers))\Rightarrow((v1_relat_1 X0)\wedge(v3_valued_0 X0)) \quad (18)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (19)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_euclid X0))\Rightarrow(v3_card_1 X1 X0)) \quad (20)$$

Theorem 1

$$\begin{aligned} &\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(m2_finseq_2 X1 k1_numbers \\ &(k1_euclid X0))\Rightarrow(\forall X2.(m2_finseq_2 X2 k1_numbers (k1_euclid \\ &X0))\Rightarrow(\forall X3.(m2_finseq_2 X3 k1_numbers (k1_euclid X0))\Rightarrow \\ &(k23_rvsum_1 (k7_euclid X0 X1 X2) X3 = k7_real_1 (k23_rvsum_1 X1 \\ &X3) (k23_rvsum_1 X2 X3)))))) \end{aligned}$$