

t20_facirc_1

(TMYtf5pQ9sp9bkYGK2HZMuQqRffWmdXX19A)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_msualg_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k2_circcomb : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $l5_struct_0 : \iota \Rightarrow o$ be given. Let $v1_msualg_1 : \iota \Rightarrow o$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u4_struct_0 : \iota \Rightarrow \iota$ be given. Let $u1_msualg_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u2_msualg_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (1)$$

Assume the following.

$$\forall X0. ((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\neg v1_xboole_0 (u1_struct_0 X0)) \quad (2)$$

Assume the following.

$$\forall X0. (l5_struct_0 X0) \Rightarrow (l1_struct_0 X0) \quad (3)$$

Assume the following.

$$\forall X0. (l1_msualg_1 X0) \Rightarrow (l5_struct_0 X0) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((\neg v2_struct_0 X0) \wedge (l1_msualg_1 X0)) \wedge \\ & ((\neg v2_struct_0 X1) \wedge (l1_msualg_1 X1))) \Rightarrow ((\neg v2_struct_0 (k2_circcomb \\ & X0 X1)) \wedge ((v1_msualg_1 (k2_circcomb X0 X1)) \wedge (l1_msualg_1 (k2_circcomb \\ & X0 X1)))) \quad (5) \end{aligned}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k2_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \vee (X3 \in X1))) \quad (6)$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge (l1_msualg_1 X0)) \Rightarrow (\forall X1. \\
& ((\neg v2_struct_0 X1) \wedge (l1_msualg_1 X1)) \Rightarrow (\forall X2.((\neg v2_struct_0 \\
& X2) \wedge ((v1_msualg_1 X2) \wedge (l1_msualg_1 X2)))) \Rightarrow ((X2 = k2_circcomb \\
& X0 X1) \Leftrightarrow ((u1_struct_0 X2 = k2_xboole_0 (u1_struct_0 X0) (u1_struct_0 \\
& X1)) \wedge ((u4_struct_0 X2 = k2_xboole_0 (u4_struct_0 X0) (u4_struct_0 \\
& X1)) \wedge ((u1_msualg_1 X2 = k1_funct_4 (u1_msualg_1 X0) (u1_msualg_1 \\
& X1)) \wedge (u2_msualg_1 X2 = k1_funct_4 (u2_msualg_1 X0) (u2_msualg_1 \\
& X1)))))))))
\end{aligned} \tag{7}$$

Assume the following.

$$\forall X0. \forall X1. k2_xboole_0 X0 X1 = k2_xboole_0 X1 X0 \tag{8}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge (l1_msualg_1 X0)) \Rightarrow (\forall X1. \\
& ((\neg v2_struct_0 X1) \wedge (l1_msualg_1 X1)) \Rightarrow (\forall X2.(m1_subset_1 \\
& X2 (u1_struct_0 X0)) \Rightarrow ((X2 \in u1_struct_0 (k2_circcomb X0 X1)) \wedge (\\
& X2 \in u1_struct_0 (k2_circcomb X1 X0))))))
\end{aligned}$$