

t20_fcont_3 (TMKAar- RAyJ2NL2VipDfrBnnapHXKcW59W2H)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v2_funct_1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarSKI : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_limfunct_1 : \iota \Rightarrow \iota$ be given. Let $k1_reset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_valued_0 : \iota \Rightarrow o$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v6_valued_0 : \iota \Rightarrow o$ be given. Let $v1_fcont_1 : \iota \Rightarrow o$ be given. Let $k2_partfun2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_reset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_funct_1 : \iota \Rightarrow \iota$ be given. Let $v1_rfunct_2 : \iota \Rightarrow o$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v2_membered : \iota \Rightarrow o$ be given. Let $v8_valued_0 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_valued_0 : \iota \Rightarrow o$ be given. Let $v7_valued_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v2_funct_1 X1) \wedge (m1_subset_1 \\ & X1 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers)))))) \Rightarrow ((v5_valued_0 \\ & (k2_partfun1 k1_numbers k1_numbers X1 X0)) \Rightarrow (v5_valued_0 (k2_partfun1 \\ & k1_numbers k1_numbers (k2_partfun2 k1_numbers k1_numbers (k2_partfun1 \\ & k1_numbers k1_numbers X1 X0)) (k7_reset_1 k1_numbers k1_numbers \\ & X1 X0)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k9_xtuple_0 (k5_relat_1 \\ & X1 X0) = k3_xboole_0 (k9_xtuple_0 X1) X0) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v2_funct_1 X0) \Rightarrow \\ & ((k10_xtuple_0 X0 = k9_xtuple_0 (k2_funct_1 X0)) \wedge (k9_xtuple_0 \\ & X0 = k10_xtuple_0 (k2_funct_1 X0)))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow ((\forall X3. (m1_subset_1 \\ & X3 X0) \Rightarrow ((X3 \in X1) \Rightarrow (X3 \in X2))) \Rightarrow (r1_tarski X1 X2))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Rightarrow (k3_xboole_0 X0 X1 = X0) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 k1_numbers k1_numbers)))) \Rightarrow (((r1_tarski X0 (k1_relset_1 \\ & k1_numbers X1)) \wedge (v1_rfunct_2 (k2_partfun1 k1_numbers k1_numbers \\ & X1 X0))) \Rightarrow ((\forall X2. (m1_subset_1 X2 k1_numbers) \Rightarrow (k7_relset_1 \\ & k1_numbers k1_numbers X1 X0 \neq k1_limfunc1 X2)) \vee (v1_fcont_1 (k2_partfun1 \\ & k1_numbers k1_numbers X1 X0)))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k10_xtuple_0 (k5_relat_1 X1 X0) = k7_relat_1 X1 X0) \quad (7)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (k7_relat_1 X0 (k9_xtuple_0 X0) = k10_xtuple_0 X0) \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v2_funct_1 X1) \wedge (m1_subset_1 \\ & X1 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers k1_numbers)))))) \Rightarrow ((v6_valued_0 \\ & (k2_partfun1 k1_numbers k1_numbers X1 X0)) \Rightarrow (v6_valued_0 (k2_partfun1 \\ & k1_numbers k1_numbers (k2_partfun2 k1_numbers k1_numbers (k2_partfun1 \\ & k1_numbers k1_numbers X1 X0)) (k7_relset_1 k1_numbers k1_numbers \\ & X1 X0)))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (k7_relset_1 X0 X1 X2 X3 = k7_relat_1 X2 X3) \quad (10)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_funct_1 X2) \wedge ((v2_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow (k2_partfun2 X0 X1 X2 = k2_funct_1 X2) \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow(k2_partfun1 \\ & X0 X1 X2 X3 = k5_relat_1 X2 X3) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(\\ & k1_relset_1 X0 X1 = k9_xtuple_0 X1) \end{aligned} \quad (13)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0)\Rightarrow(k5_relat_1 X0 (k9_xtuple_0 X0) = X0) \quad (14)$$

Assume the following.

$$v3_membered\ k1_numbers \quad (15)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v2_funct_1 \\ & X1)))\Rightarrow((v1_relat_1 (k5_relat_1 X1 X0))\wedge(v2_funct_1 (k5_relat_1 \\ & X1 X0))) \end{aligned} \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 X0)\Rightarrow(v1_relat_1 (k5_relat_1 X0 X1)) \quad (17)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((v1_funct_1 X2)\wedge((v2_funct_1 \\ & X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((v1_funct_1 \\ & (k2_partfun2 X0 X1 X2))\wedge(m1_subset_1 (k2_partfun2 X0 X1 X2) (k1_zfmisc_1 \\ & (k2_zfmisc_1 X1 X0)))) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((v1_funct_1 \\ & (k2_partfun1 X0 X1 X2 X3))\wedge(m1_subset_1 (k2_partfun1 X0 X1 X2 X3) \\ & (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \end{aligned} \quad (19)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow((v1_relat_1 (k2_funct_1 X0))\wedge(v1_funct_1 (k2_funct_1 X0))) \quad (20)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(m1_subset_1 (k1_relset_1 X0 X1) (k1_zfmisc_1 X0)) \quad (21)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (22)$$

Assume the following.

$$\forall X0.(v3_membered X0)\Rightarrow(v2_membered X0) \quad (23)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers k1_numbers)))\Rightarrow(((v1_funct_1 X0)\wedge(v8_valued_0 X0))\Rightarrow((v1_funct_1 X0)\wedge(v1_rfunct_2 X0))) \quad (24)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (25)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v2_valued_0 X0)\wedge(v6_valued_0 X0))))\Rightarrow((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v2_valued_0 X0)\wedge(v8_valued_0 X0)))) \quad (26)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v2_valued_0 X0)\wedge(v5_valued_0 X0))))\Rightarrow((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v2_valued_0 X0)\wedge(v7_valued_0 X0)))) \quad (27)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers k1_numbers)))\Rightarrow(((v1_funct_1 X0)\wedge(v7_valued_0 X0))\Rightarrow((v1_funct_1 X0)\wedge(v1_rfunct_2 X0))) \quad (28)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (29)$$

Assume the following.

$$\forall X0.\forall X1.(v2_membered X1)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v2_valued_0 X2)) \quad (30)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (\forall X1.((v1_funct_1 \\ & X1) \wedge ((v2_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k1_numbers k1_numbers)))))) \Rightarrow ((r1_tarski (k1_limfunc1 X0) (k1_relset_1 \\ & k1_numbers X1)) \Rightarrow (((\neg v5_valued_0 (k2_partfun1 k1_numbers k1_numbers \\ & X1 (k1_limfunc1 X0))) \wedge (\neg v6_valued_0 (k2_partfun1 k1_numbers \\ & k1_numbers X1 (k1_limfunc1 X0)))) \vee (v1_fcont_1 (k2_partfun1 k1_numbers \\ & k1_numbers (k2_partfun2 k1_numbers k1_numbers (k2_partfun1 k1_numbers \\ & k1_numbers X1 (k1_limfunc1 X0))) (k7_relset_1 k1_numbers k1_numbers \\ & X1 (k1_limfunc1 X0)))))) \end{aligned}$$