

t20_fdifff_3 (TMF- pGu4Bvk2JRTuJ4Wi3wn4LTkdb5WCTcRr)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r3_fdiff_3 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k6_rfunct_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_fdiff_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $k10_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_square_1 : \iota \Rightarrow \iota$ be given. Let $r2_fdiff_3 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_rcomp_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v2_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_fdiff_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_funct_1 : \iota \Rightarrow o$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k3_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_comseq_2 : \iota \Rightarrow o$ be given. Let $k20_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k37_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k47_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
& \forall X0.((v1_funct_1 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\
& k1_numbers k1_numbers)))) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow \\
& (\forall X2.(m1_subset_1 X2 k1_numbers) \Rightarrow (\neg(r2_fdiff_3 X0 X1) \wedge \\
& ((k1_seq_1 X0 X1 \neq X2) \wedge ((\exists X3.(m1_subset_1 X3 k1_numbers) \wedge \\
& ((\neg r1_xxreal_0 X3 k6_numbers) \wedge (r1_tarski (k1_rcomp_1 X1 (k7_real_1 \\
& X1 X3)) (k1_relset_1 k1_numbers X0)))) \wedge (\forall X3.(m1_subset_1 \\
& X3 k1_numbers) \Rightarrow (\neg(\neg r1_xxreal_0 X3 k6_numbers) \wedge ((r1_tarski (\\
& k1_rcomp_1 X1 (k7_real_1 X1 X3)) (k1_relset_1 k1_numbers X0)) \wedge \\
& (\forall X4.(m1_subset_1 X4 k1_numbers) \Rightarrow (\neg(X4 \in k1_rcomp_1 X1 \\
& (k7_real_1 X1 X3)) \wedge (k1_seq_1 X0 X4 = X2))))))))))
\end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_funct_1 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ k1_numbers k1_numbers)))) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow \\ ((r3_fdiff_3 X0 X1) \Rightarrow (r2_fdiff_3 X0 X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_funct_1 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ k1_numbers k1_numbers)))) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow \\ ((r3_fdiff_3 X0 X1) \Rightarrow ((\forall X2.(m1_subset_1 X2 k1_numbers) \Rightarrow \\ (\neg(\neg r1_xxreal_0 X2 k6_numbers) \wedge (\forall X3.(m1_subset_1 X3 k1_numbers) \Rightarrow \\ (\neg(X3 \in k1_relset_1 k1_numbers X0) \wedge ((X3 \in k1_rcomp_1 X1 (k7_real_1 \\ X1 X2)) \wedge (k1_seq_1 X0 X3 = k6_numbers)))))) \vee ((r3_fdiff_3 (k6_rfunct_1 \\ k1_numbers k1_numbers X0) X1) \wedge (k2_fdiff_3 X1 (k6_rfunct_1 k1_numbers \\ k1_numbers X0) = k1_real_1 (k10_real_1 (k2_fdiff_3 X1 X0) (k5_square_1 \\ (k1_seq_1 X0 X1)))))))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 X0))) \Rightarrow (m1_subset_1 (k1_seq_1 X0 X1) k1_numbers) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_funct_1 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ k1_numbers k1_numbers)))) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow \\ ((r3_fdiff_3 X0 X1) \Leftrightarrow ((\exists X2.(m1_subset_1 X2 k1_numbers) \wedge \\ ((\neg r1_xxreal_0 X2 k6_numbers) \wedge (r1_tarski (k1_rcomp_1 X1 (k7_real_1 \\ X1 X2)) (k1_relset_1 k1_numbers X0)))) \wedge (\forall X2.((v2_relat_1 \\ X2) \wedge ((v1_funct_1 X2) \wedge ((v1_funct_2 X2 k5_numbers k1_numbers) \wedge \\ ((v1_fdiff_1 X2 k6_numbers) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ k5_numbers k1_numbers)))))) \Rightarrow (\forall X3.((v1_funct_1 X3) \wedge \\ ((v3_funct_1 X3) \wedge ((v1_funct_2 X3 k5_numbers k1_numbers) \wedge (m1_subset_1 \\ X3 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow ((\\ (k2_relset_1 k1_numbers X3 = k1_tarski X1) \wedge ((r1_tarski (k2_relset_1 \\ k1_numbers (k3_valued_1 k5_numbers k1_numbers k1_numbers X2 X3)) \\ (k1_relset_1 k1_numbers X0)) \wedge (\forall X4.(m2_subset_1 X4 k1_numbers \\ k5_numbers) \Rightarrow (\neg r1_xxreal_0 (k1_seq_1 X2 X4) k6_numbers)))) \Rightarrow (\\ v2_comseq_2 (k20_valued_1 k5_numbers k1_numbers k1_numbers (\\ k37_valued_1 k5_numbers k1_numbers X2) (k47_valued_1 k5_numbers \\ k1_numbers k1_numbers (k8_funct_2 k5_numbers k1_numbers k1_numbers \\ (k3_valued_1 k5_numbers k1_numbers k1_numbers X2 X3) X0) (k8_funct_2 \\ k5_numbers k1_numbers k1_numbers X3 X0)))))))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v5_relat_1 X0 k1_numbers)) \Rightarrow ((v1_relat_1 X0) \wedge (v3_valued_0 X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (8)$$

Theorem 1

$$\begin{aligned} &\forall X0.((v1_funct_1 X0)\wedge(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ &\quad k1_numbers k1_numbers))))\Rightarrow(\forall X1.(m1_subset_1 X1 k1_numbers)\Rightarrow \\ &((r3_fdiff_3 X0 X1)\Rightarrow((k1_seq_1 X0 X1 = k6_numbers)\vee((r3_fdiff_3 \\ &\quad (k6_rfunct_1 k1_numbers k1_numbers X0) X1)\wedge(k2_fdiff_3 X1 (k6_rfunct_1 \\ &\quad k1_numbers k1_numbers X0) = k1_real_1 (k10_real_1 (k2_fdiff_3 \\ &\quad X1 X0) (k5_square_1 (k1_seq_1 X0 X1)))))))) \end{aligned}$$