

# t20\_fintopo3 (TMRgUZHpBTByiBrMx- aKaMekQZprVCWnRkjJ)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $l1\_orders\_2 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k5\_fintopo3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k8\_fin\_topo : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $np\_0 : \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k4\_fintopo3 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$(m2\_subset\_1 np\_0 k1\_numbers k5\_numbers) \wedge ((m1\_subset\_1 np\_0 k5\_numbers) \wedge (m1\_subset\_1 np\_0 k1\_numbers)) \quad (3)$$

Assume the following.

$$v1\_xboole\_0 np\_0 \quad (4)$$

Assume the following.

$$k2\_xcmplx\_0 np\_0 np\_1 = np\_1 \quad (5)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (6)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((\neg v1\_xboole\_0 X0) \wedge \\ & (((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1)))))) \wedge (m1\_subset\_1 X3 X0))) \Rightarrow (k3\_funct\_2 X0 \\ & X1 X2 X3 = k1\_funct\_1 X2 X3) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((m1\_subset\_1 X0 k5\_numbers) \wedge (v7\_ordinal1 \\ & X1)) \Rightarrow (k2\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \end{aligned} \quad (9)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1) \wedge (v3\_ordinal1 k4\_ordinal1) \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((\neg v2\_struct\_0 X0) \wedge (l1\_orders\_2 X0)) \wedge \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0)))) \Rightarrow ((v1\_funct\_1 \\ & (k4\_fintopo3 X0 X1)) \wedge ((v1\_funct\_2 (k4\_fintopo3 X0 X1) k5\_numbers \\ & (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \wedge (m1\_subset\_1 (k4\_fintopo3 X0 \\ & X1) (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers (k1\_zfmisc\_1 (u1\_struct\_0 \\ & X0)))))) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2\_struct\_0 X0) \wedge (l1\_orders\_2 X0)) \Rightarrow (\forall X1. \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow (\forall X2. \\ & (v7\_ordinal1 X2) \Rightarrow (k5\_fintopo3 X0 X1 X2 = k1\_funct\_1 (k4\_fintopo3 \\ & X0 X1) X2))) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2\_struct\_0 X0) \wedge (l1\_orders\_2 X0)) \Rightarrow (\forall X1. \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow (\forall X2. \\ & ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 X2 k5\_numbers (k1\_zfmisc\_1 (u1\_struct\_0 \\ & X0))) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers ( \\ & k1\_zfmisc\_1 (u1\_struct\_0 X0)))))) \Rightarrow ((X2 = k4\_fintopo3 X0 X1) \Leftrightarrow \\ & ((\forall X3. (m1\_subset\_1 X3 k5\_numbers) \Rightarrow (\forall X4. (m1\_subset\_1 \\ & X4 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow ((X4 = k3\_funct\_2 k5\_numbers \\ & (k1\_zfmisc\_1 (u1\_struct\_0 X0)) X2 X3) \Rightarrow (k3\_funct\_2 k5\_numbers \\ & (k1\_zfmisc\_1 (u1\_struct\_0 X0)) X2 (k2\_nat\_1 X3 np\_1) = k8\_fin\_topo \\ & X0 X4)))) \wedge (k3\_funct\_2 k5\_numbers (k1\_zfmisc\_1 (u1\_struct\_0 X0)) \\ & X2 k6\_numbers = X1)))))) \end{aligned} \quad (13)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (14)$$

**Theorem 1**

$$\begin{aligned} \forall X0.((\neg v2\_struct\_0 X0) \wedge (l1\_orders\_2 X0)) \Rightarrow (\forall X1. \\ (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow (k5\_fintopo3 \\ X0 X1 np\_1 = k8\_fin\_topo X0 X1)) \end{aligned}$$