

t20_funct_2

(TMEo57V21usTHGnNzaiVXBZyKqXCH2GSWZU)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_funct_1 : \iota \Rightarrow o$ be given. Let $k1_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarSKI : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (((v2_funct_1 (k3_relat_1 X1 X0)) \wedge (r1_tarSKI (k10_xtuple_0 X1) (k9_xtuple_0 X0))) \Rightarrow (v2_funct_1 X1))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. \forall X4. \forall X5. (((v1_funct_1 X4) \wedge (m1_subset_1 X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \wedge ((v1_funct_1 X5) \wedge (m1_subset_1 X5 (k1_zfmisc_1 (k2_zfmisc_1 X2 X3)))))) \Rightarrow (k1_partfun1 X0 X1 X2 X3 X4 X5 = k3_relat_1 X4 X5) \quad (3)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (((X1 \neq k1_xboole_0) \Rightarrow ((v1_funct_2 X2 X0 X1) \Leftrightarrow (X0 = k1_relset_1 X0 X2))) \wedge ((X1 = k1_xboole_0) \Rightarrow ((v1_funct_2 X2 X0 X1) \Leftrightarrow (X2 = k1_xboole_0)))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 X1)\Rightarrow((v5_relat_1 X1 X0)\Leftrightarrow(r1_tarski (k10_xtuple_0 X1) X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (7)$$

Assume the following.

$$\forall X0.((v1_xboole_0 X0)\wedge((v1_relat_1 X0)\wedge(v1_funct_1 X0)))\Rightarrow ((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v2_funct_1 X0))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(v1_relat_1 X0) \quad (10)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(v1_funct_1 X0) \quad (11)$$

Theorem 1

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X3)\wedge \\ & ((v1_funct_2 X3 X0 X1)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1))))\Rightarrow(\forall X4.((v1_funct_1 X4)\wedge((v1_funct_2 X4 X1 X2)\wedge \\ & (m1_subset_1 X4 (k1_zfmisc_1 (k2_zfmisc_1 X1 X2))))\Rightarrow((v2_funct_1 \\ & (k1_partfun1 X0 X1 X1 X2 X3 X4))\Rightarrow(((X2 = k1_xboole_0)\wedge(X1\neq k1_xboole_0))\vee \\ & (v2_funct_1 X3)))) \end{aligned}$$