

t20_mod_2

(TMcZhwLH6tgRvSnN9jb4dXqbHVfLJQPXpPi)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_classes2 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_ordinal1 : \iota \Rightarrow \iota$ be given. Let $k7_classes2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_classes2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (((\neg v1_xboole_0 X0) \wedge (v1_classes2 X0)) \wedge ((m1_subset_1 X1 X0) \wedge (m1_subset_1 X2 X0))) \Rightarrow (k7_classes2 X0 X1 X2 = k2_xboole_0 X1 X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (((\neg v1_xboole_0 X0) \wedge (v1_classes2 X0)) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k1_classes2 X0 X1 = k1_tarski X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (((\neg v1_xboole_0 X0) \wedge (v1_classes2 X0)) \wedge ((m1_subset_1 X1 X0) \wedge (m1_subset_1 X2 X0))) \Rightarrow (m1_subset_1 (k7_classes2 X0 X1 X2) X0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (((\neg v1_xboole_0 X0) \wedge (v1_classes2 X0)) \wedge (m1_subset_1 X1 X0)) \Rightarrow (m1_subset_1 (k1_classes2 X0 X1) X0) \quad (4)$$

Assume the following.

$$\forall X0. k1_ordinal1 X0 = k2_xboole_0 X0 (k1_tarski X0) \quad (5)$$

Theorem 1

$$\forall X0. ((\neg v1_xboole_0 X0) \wedge (v1_classes2 X0)) \Rightarrow (\forall X1. (m1_subset_1 X1 X0) \Rightarrow (m1_subset_1 (k1_ordinal1 X1) X0))$$