

t20_pepin (TMEpgKDJYBptPBkyuBd- EqF7hAMdvfXqYtZd)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_abian : \iota \Rightarrow o$ be given. Let $k1_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_0 : \iota$ be given. Let $v1_int_1 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow (k1_newton X1 (k1_nat_1 X0 np_1) = k3_xcmplx_0 (k1_newton X1 X0) X1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k1_newton X0 k6_numbers = np_1) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 np_1 X0 = X0) \quad (4)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\neg (\neg v1_abian X0) \wedge ((\neg v1_abian X1) \wedge (v1_abian (k3_xcmplx_0 X0 X1)))))) \quad (5)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (6)$$

Assume the following.

$$\forall X0 : \iota \Rightarrow o. ((X0 k6_numbers) \wedge (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X0 X1) \Rightarrow (X0 (k1_nat_1 X1 np_1)))))) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (X0 X1)) \quad (7)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_int_1 X0)\wedge(v1_abian X0))\wedge(v1_int_1 X1))\Rightarrow(v1_abian (k3_xcmplx_0 X0 X1)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(v7_ordinal1 (k1_newton X0 X1)) \quad (10)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(v7_ordinal1 X0) \quad (11)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xcmplx_0 X0) \quad (12)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_xreal_0 X0) \quad (13)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_int_1 X0) \quad (14)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(v7_ordinal1 X1)\Rightarrow(\neg(\neg v1_abian X0)\wedge(v1_abian (k1_newton X0 X1))))$$