

t20_prob_2
(TMcWU1iESr27pdybCxtFSNsQysgTsky1EAW)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m1_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r2_prob_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_prob_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
& \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2. (m2_prob_1 X2 X0 X1) \Rightarrow (\forall X3. \\
& (m1_prob_1 X3 X0 X1) \Rightarrow (\forall X4. (m1_prob_1 X4 X0 X1) \Rightarrow (\forall X5. \\
& (m1_prob_1 X5 X0 X1) \Rightarrow ((r2_prob_2 X0 X1 X2 X3 X4 X5) \Leftrightarrow ((k1_seq_1 X2 \\
& (k5_prob_1 X0 X1 (k5_prob_1 X0 X1 X3 X4) X5) = k8_real_1 (k8_real_1 \\
& (k1_seq_1 X2 X3) (k1_seq_1 X2 X4)) (k1_seq_1 X2 X5)) \wedge ((k1_seq_1 \\
& X2 (k5_prob_1 X0 X1 X3 X4) = k8_real_1 (k1_seq_1 X2 X3) (k1_seq_1 X2 \\
& X4)) \wedge ((k1_seq_1 X2 (k5_prob_1 X0 X1 X3 X5) = k8_real_1 (k1_seq_1 \\
& X2 X3) (k1_seq_1 X2 X5)) \wedge (k1_seq_1 X2 (k5_prob_1 X0 X1 X4 X5) = k8_real_1 \\
& (k1_seq_1 X2 X4) (k1_seq_1 X2 X5))))))))))
\end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
& \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2. (m2_prob_1 X2 X0 X1) \Rightarrow (\forall X3. \\
& (m1_prob_1 X3 X0 X1) \Rightarrow (\forall X4. (m1_prob_1 X4 X0 X1) \Rightarrow ((r1_prob_2 \\
& X0 X1 X2 X3 X4) \Leftrightarrow (k1_seq_1 X2 (k5_prob_1 X0 X1 X3 X4) = k8_real_1 (k1_seq_1 \\
& X2 X3) (k1_seq_1 X2 X4))))))
\end{aligned} \tag{2}$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\ & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.(m1_prob_1 X2 X0 X1) \Rightarrow (\forall X3. \\ & (m1_prob_1 X3 X0 X1) \Rightarrow (\forall X4.(m1_prob_1 X4 X0 X1) \Rightarrow (\forall X5. \\ & (m2_prob_1 X5 X0 X1) \Rightarrow ((r2_prob_2 X0 X1 X5 X2 X3 X4) \Leftrightarrow ((k1_seq_1 X5 \\ & (k5_prob_1 X0 X1 (k5_prob_1 X0 X1 X2 X3) X4) = k8_real_1 (k8_real_1 \\ & (k1_seq_1 X5 X2) (k1_seq_1 X5 X3)) (k1_seq_1 X5 X4)) \wedge ((r1_prob_2 \\ & X0 X1 X5 X2 X3) \wedge ((r1_prob_2 X0 X1 X5 X3 X4) \wedge (r1_prob_2 X0 X1 X5 X2 X4)))))))))) \end{aligned}$$