

t20_tdlat_1

(TMdKy2M8ymMvnzG9pFpXsL4fejQRdPUQ2X4)

October 27, 2020

Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $v6_tops_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v4_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tops_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(l1_pre_topc X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ (u1_struct_0 X0))) \Rightarrow (((v4_pre_topc X1 X0) \Rightarrow (k2_pre_topc X0 X1 = \\ X1)) \wedge (((v2_pre_topc X0) \wedge (k2_pre_topc X0 X1 = X1)) \Rightarrow (v4_pre_topc \\ X1 X0)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.\exists X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \wedge (v1_xboole_0 X1) \quad (3)$$

Assume the following.

$$\forall X0.(l1_pre_topc X0) \Rightarrow (v1_xboole_0 (k1_tops_1 X0 (k1_struct_0 X0))) \quad (4)$$

Assume the following.

$$\forall X0.(l1_pre_topc X0) \Rightarrow (l1_struct_0 X0) \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.(l1_pre_topc X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ (u1_struct_0 X0))) \Rightarrow ((v6_tops_1 X1 X0) \Leftrightarrow (X1 = k1_tops_1 X0 (k2_pre_topc \\ X0 X1)))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(l1_struct_0 X0) \Rightarrow (k1_struct_0 X0 = k1_xboole_0) \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. \\ (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((v1_xboole_0 \\ X1) \Rightarrow (v4_pre_topc X1 X0))) \end{aligned} \quad (8)$$

Theorem 1

$$\forall X0.((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow (v6_tops_1 (k1_struct_0 X0) X0)$$