

t20_tmap_1 (TM- PHng9hvmBB3bAQxNp1d7K3PTskpjaDTxj)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_tsep_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $g1_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $u1_pre_topc : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(l1_pre_topc X0) \Rightarrow (m1_pre_topc X0 X0) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc \\ & X0))) \Rightarrow (\forall X1.((\neg v2_struct_0 X1) \wedge (m1_pre_topc X1 X0)) \Rightarrow (\\ & \forall X2.((\neg v2_struct_0 X2) \wedge (m1_pre_topc X2 X0)) \Rightarrow ((m1_pre_topc \\ & X1 X2) \Leftrightarrow (k1_tsep_1 X0 X1 X2 = g1_pre_topc (u1_struct_0 X2) (u1_pre_topc \\ & X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.(l1_pre_topc X0) \Rightarrow (\forall X1.(m1_pre_topc X1 X0) \Rightarrow (l1_pre_topc X1)) \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc \\ & X0))) \Rightarrow (\forall X1.((\neg v2_struct_0 X1) \wedge (m1_pre_topc X1 X0)) \Rightarrow (\\ & k1_tsep_1 X0 X1 X1 = g1_pre_topc (u1_struct_0 X1) (u1_pre_topc X1))) \end{aligned}$$