

t20_toprealc (TMawiRLCdkWetSGDEPUkMFB-
ByfSVTSJSYP3)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k12_euclid : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_rvsum_1 : \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_square_1 : \iota \Rightarrow \iota$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k39_valued_1 : \iota \Rightarrow \iota$ be given. Let $k3_square_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k12_rvsum_1 : \iota \Rightarrow \iota$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 k1_numbers) \Rightarrow ((\forall X1.(m2_subset_1 \\ & X1 k1_numbers k5_numbers) \Rightarrow ((X1 \in k4_finseq_1 X0) \Rightarrow (k1_seq_1 X0 \\ & X1 = k6_numbers))) \Rightarrow (k18_rvsum_1 X0 = k6_numbers)) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 X0 k6_numbers = k6_numbers) \quad (2)$$

Assume the following.

$$k7_square_1 k6_numbers = k6_numbers \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ & ((k9_xtuple_0 (k39_valued_1 X0) = k9_xtuple_0 X0) \wedge (\forall X1. \\ & k1_funct_1 (k39_valued_1 X0) X1 = k3_square_1 (k1_funct_1 X0 X1))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Leftrightarrow(m1_subset_1 X2 X1)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Leftrightarrow(m1_finseq_1 X1 X0) \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow(k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0)))\Rightarrow(k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (9)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v3_valued_0 X0)\wedge(v1_finseq_1 X0))))\Rightarrow(k12_rvsum_1 X0 = k39_valued_1 X0) \quad (10)$$

Assume the following.

$$\neg v1_finset_1 k4_ordinal1 \quad (11)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (12)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Rightarrow(m1_subset_1 X2 X0)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge(v1_funct_1 X1)\wedge(v1_finseq_1 X1)) \quad (16)$$

Assume the following.

$$m2_subset_1 k6_numbers k1_numbers k5_numbers \quad (17)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (18)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v3_valued_0 X0)\wedge(v1_finseq_1 X0))))\Rightarrow(m2_finseq_1 (k12_rvsum_1 X0) k1_numbers) \quad (19)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v1_finseq_1 X0)\wedge(v3_valued_0 X0))))\Rightarrow(k12_euclid X0 = k7_square_1 (k18_rvsum_1 (k12_rvsum_1 X0))) \quad (20)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0)\Rightarrow(k3_square_1 X0 = k3_xcmplx_0 X0 X0) \quad (21)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v3_valued_0 X0))\Rightarrow((v1_relat_1 X0)\wedge(v1_valued_0 X0)) \quad (22)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow(v5_relat_1 X1 X0) \quad (23)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v5_relat_1 X0 k1_numbers))\Rightarrow((v1_relat_1 X0)\wedge(v3_valued_0 X0)) \quad (24)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xcmplx_0 X0) \quad (25)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(v1_finset_1 X0) \quad (26)$$

Assume the following.

$$\forall X0.(v6_membered X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(v7_ordinal1 X1)) \quad (27)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (\neg(k12_euclid X0 \neq k6_numbers) \wedge (\forall X1. (v7_ordinal1 X1) \Rightarrow (\neg(X1 \in k4_finseq_1 X0) \wedge (k1_funct_1 X0 X1 \neq k6_numbers))))))$$