

t20_waybel_0
(TMYy7G9PTv5cDYdNcRGdcj3qiaPKLXkj1Uc)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v3_orders_2 : \iota \Rightarrow o$ be given. Let $v5_orders_2 : \iota \Rightarrow o$ be given. Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k6_waybel_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_orders_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. ((\neg v2_struct_0 X0) \wedge (l1_orders_2 X0)) \Rightarrow (\forall X1. \\ (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. (m1_subset_1 X2 \\ (u1_struct_0 X0)) \Rightarrow ((X2 \in k6_waybel_0 X0 X1) \Leftrightarrow (r1_orders_2 X0 X1 \\ X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0. (l1_orders_2 X0) \Rightarrow ((v5_orders_2 X0) \Leftrightarrow (\forall X1. (\\ m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. (m1_subset_1 X2 \\ (u1_struct_0 X0)) \Rightarrow ((r1_orders_2 X0 X1 X2) \wedge (r1_orders_2 X0 X2 \\ X1)) \Rightarrow (X1 = X2)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0. ((\neg v2_struct_0 X0) \wedge (l1_orders_2 X0)) \Rightarrow ((v3_orders_2 \\ X0) \Leftrightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (r1_orders_2 \\ X0 X1 X1))) \end{aligned} \tag{3}$$

Theorem 1

$$\begin{aligned} \forall X0. ((\neg v2_struct_0 X0) \wedge ((v3_orders_2 X0) \wedge ((v5_orders_2 \\ X0) \wedge (l1_orders_2 X0)))) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 \\ X0)) \Rightarrow (\forall X2. (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow ((k6_waybel_0 \\ X0 X1 = k6_waybel_0 X0 X2) \Rightarrow (X1 = X2)))) \end{aligned}$$