

# t20\_xboolean (TMT- PdWidWc6oj4Z2kYfRwJXeGMSGEUFXNmB)

October 27, 2020

Let  $v1\_xboolean : \iota \Rightarrow o$  be given. Let  $k4\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $np\_0 : \iota$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xboolean : \iota \Rightarrow \iota$  be given. Let  $k2\_xboolean : \iota$  be given. Let  $k1\_xboolean : \iota$  be given. Let  $k6\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X_0. (v1\_xboole\_0 X_0) \Rightarrow (X_0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\forall X_0. (v1\_xcmplx\_0 X_0) \Rightarrow (k3\_xcmplx\_0 np\_1 X_0 = X_0) \quad (2)$$

Assume the following.

$$\forall X_0. (v1\_xcmplx\_0 X_0) \Rightarrow (k3\_xcmplx\_0 X_0 k6\_numbers = k6\_numbers) \quad (3)$$

Assume the following.

$$v1\_xboole\_0 np\_0 \quad (4)$$

Assume the following.

$$k6\_xcmplx\_0 np\_1 np\_1 = np\_0 \quad (5)$$

Assume the following.

$$k6\_xcmplx\_0 np\_1 np\_0 = np\_1 \quad (6)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (7)$$

Assume the following.

$$\forall X_0. (v1\_xboolean X_0) \Rightarrow (k3\_xboolean (k3\_xboolean X_0) = X_0) \quad (8)$$

Assume the following.

$$\forall X_0 \forall X_1. ((v1\_xboolean X_0) \wedge (v1\_xboolean X_1)) \Rightarrow (k4\_xboolean X_0 X_0 = X_0) \quad (9)$$

Assume the following.

$$v1\_xboolean k2\_xboolean \quad (10)$$

Assume the following.

$$v1\_xboolean k1\_xboolean \quad (11)$$

Assume the following.

$$\forall X_0. (v1\_xboolean X_0) \Rightarrow (v1\_xboolean (k3\_xboolean X_0)) \quad (12)$$

Assume the following.

$$\forall X_0. (v1\_xboolean X_0) \Rightarrow (\forall X_1. (v1\_xboolean X_1) \Rightarrow (k7\_xboolean X_0 X_1 = k4\_xboolean (k6\_xboolean X_0 X_1) (k6\_xboolean X_1 X_0))) \quad (13)$$

Assume the following.

$$\forall X_0. (v1\_xboolean X_0) \Rightarrow (\forall X_1. (v1\_xboolean X_1) \Rightarrow (k6\_xboolean X_0 X_1 = k5\_xboolean (k3\_xboolean X_0) X_1)) \quad (14)$$

Assume the following.

$$\forall X_0. (v1\_xboolean X_0) \Rightarrow (\forall X_1. (v1\_xboolean X_1) \Rightarrow (k5\_xboolean X_0 X_1 = k3\_xboolean (k4\_xboolean (k3\_xboolean X_0) (k3\_xboolean X_1)))) \quad (15)$$

Assume the following.

$$\forall X_0. (v1\_xboolean X_0) \Rightarrow (\forall X_1. (v1\_xboolean X_1) \Rightarrow (k4\_xboolean X_0 X_1 = k3\_xcmplx_0 X_0 X_1)) \quad (16)$$

Assume the following.

$$\forall X_0. (v1\_xboolean X_0) \Rightarrow (k3\_xboolean X_0 = k6\_xcmplx_0 np\_1 X_0) \quad (17)$$

Assume the following.

$$\forall X_0. (v1\_xboolean X_0) \Leftrightarrow ((X_0 = k1\_xboolean) \vee (X_0 = k2\_xboolean)) \quad (18)$$

Assume the following.

$$k2\_xboolean = np\_1 \quad (19)$$

Assume the following.

$$k1\_xboolean = k6\_numbers \quad (20)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_xboolean\ X0) \wedge (v1\_xboolean\ X1)) \Rightarrow (k5\_xboolean\ X0\ X1 = k5\_xboolean\ X1\ X0) \quad (21)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_xboolean\ X0) \wedge (v1\_xboolean\ X1)) \Rightarrow (k4\_xboolean\ X0\ X1 = k4\_xboolean\ X1\ X0) \quad (22)$$

Assume the following.

$$\forall X0. (v7\_ordinal1\ X0) \Rightarrow (v1\_xcmplx_0\ X0) \quad (23)$$

Assume the following.

$$\forall X0. (v1\_xboolean\ X0) \Rightarrow (v7\_ordinal1\ X0) \quad (24)$$

### Theorem 1

$$\forall X0. (v1\_xboolean\ X0) \Rightarrow (\forall X1. (v1\_xboolean\ X1) \Rightarrow (k4\_xboolean\ X0\ (k7\_xboolean\ X0\ X1) = k4\_xboolean\ X0\ X1)))$$