

t20_zf_model (TMJB-
VkVFu6iz7oaFf8Wtyxd9fP3HDnXVWKJ)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zf_lang : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_zf_lang : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_zf_model : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k13_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_zf_lang : \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\
 & (v1_funct_2 X1 k1_zf_lang X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\
 & k1_zf_lang X0)))))) \Rightarrow (\forall X2.((v1_zf_lang X2) \wedge (m2_finseq_1 \\
 & X2 k5_numbers)) \Rightarrow (\forall X3.(m2_subset_1 X3 k5_numbers k1_zf_lang) \Rightarrow \\
 & ((r1_zf_model X0 X1 (k8_zf_lang X3 X2)) \Leftrightarrow (\forall X4.((v1_funct_1 \\
 & X4) \wedge ((v1_funct_2 X4 k1_zf_lang X0) \wedge (m1_subset_1 X4 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 k1_zf_lang X0)))))) \Rightarrow ((\forall X5.(m2_subset_1 X5 \\
 & k5_numbers k1_zf_lang) \Rightarrow ((k3_funct_2 k1_zf_lang X0 X4 X5 \neq k3_funct_2 \\
 & k1_zf_lang X0 X1 X5) \Rightarrow (X3 = X5))) \Rightarrow (r1_zf_model X0 X4 X2))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\
 & (v1_funct_2 X1 k1_zf_lang X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\
 & k1_zf_lang X0)))))) \Rightarrow (\forall X2.((v1_zf_lang X2) \wedge (m2_finseq_1 \\
 & X2 k5_numbers)) \Rightarrow ((r1_zf_model X0 X1 X2) \Leftrightarrow (\neg r1_zf_model X0 X1 (k6_zf_lang \\
 & X2))))))
 \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\
 & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\
 & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1))
 \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Leftrightarrow(m1_finseq_1 X1 X0) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_zf_lang)\wedge((v1_zf_lang X1)\wedge(m1_finseq_1 X1 k5_numbers)))\Rightarrow(v1_zf_lang (k8_zf_lang X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0.((v1_zf_lang X0)\wedge(m1_finseq_1 X0 k5_numbers))\Rightarrow(v1_zf_lang (k6_zf_lang X0)) \quad (6)$$

Assume the following.

$$\neg v1_xboole_0 k1_zf_lang \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_zf_lang)\wedge(m1_finseq_1 X1 k5_numbers))\Rightarrow(m2_finseq_1 (k8_zf_lang X0 X1) k5_numbers) \quad (8)$$

Assume the following.

$$\forall X0.(m1_finseq_1 X0 k5_numbers)\Rightarrow(m2_finseq_1 (k6_zf_lang X0) k5_numbers) \quad (9)$$

Assume the following.

$$m1_subset_1 k1_zf_lang (k1_zfmisc_1 k5_numbers) \quad (10)$$

Assume the following.

$$\forall X0.(m2_subset_1 X0 k5_numbers k1_zf_lang)\Rightarrow(\forall X1.((v1_zf_lang X1)\wedge(m2_finseq_1 X1 k5_numbers))\Rightarrow(k13_zf_lang X0 X1 = k6_zf_lang (k8_zf_lang X0 (k6_zf_lang X1)))) \quad (11)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_xboole_0 X1)) \quad (12)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge \\ & (v1_funct_2 X1 k1_zf_lang X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k1_zf_lang X0))))\Rightarrow(\forall X2.((v1_zf_lang X2)\wedge(m2_finseq_1 \\ & X2 k5_numbers))\Rightarrow(\forall X3.(m2_subset_1 X3 k5_numbers k1_zf_lang)\Rightarrow \\ & ((r1_zf_model X0 X1 (k13_zf_lang X3 X2))\Leftrightarrow(\exists X4.((v1_funct_1 \\ & X4)\wedge((v1_funct_2 X4 k1_zf_lang X0)\wedge(m1_subset_1 X4 (k1_zfmisc_1 \\ & (k2_zfmisc_1 k1_zf_lang X0))))\wedge((\forall X5.(m2_subset_1 X5 \\ & k5_numbers k1_zf_lang)\Rightarrow((k3_funct_2 k1_zf_lang X0 X4 X5\neq k3_funct_2 \\ & k1_zf_lang X0 X1 X5)\Rightarrow(X3 = X5))))\wedge(r1_zf_model X0 X4 X2)))))) \end{aligned}$$