

t218_xxreal_1

(TMVfQ5uxRuzztSWNASmeBeqxxEBJ3wocP6k)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_0 : \iota$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (r1_xxreal_0 X0 k1_xxreal_0) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((X0 \in k3_xxreal_1 X1 X2) \Leftrightarrow ((\neg r1_xxreal_0 X0 X1) \wedge \\ & (r1_xxreal_0 X0 X2))))) \end{aligned} \quad (2)$$

Assume the following.

$$v1_xxreal_0 k1_xxreal_0 \quad (3)$$

Theorem 1

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((\neg r1_xxreal_0 X0 X1) \Rightarrow (X0 \in k3_xxreal_1 X1 k1_xxreal_0)))$$