

t21_algspec1

(TMZa5xGX7r7ATLnRrfUne2tDzJ5QfpWoFgc)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_algspec1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_partfun1 : \iota \Rightarrow \iota$ be given. Let $k4_relat_1 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge \\ (v1_funct_1 X2)) \Rightarrow (k3_relat_1 X0 (k1_funct_4 X1 X2) = k1_funct_4 \\ (k3_relat_1 X0 X1) (k3_relat_1 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.k6_partfun1 X0 = k4_relat_1 X0 \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2.(r1_tarski (k10_xtuple_0 \\ X0) X2) \Rightarrow (k3_relat_1 X0 (k5_relat_1 X1 X2) = k3_relat_1 X0 X1))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v1_relat_1 \\ (k5_relat_1 X0 X1)) \wedge (v1_funct_1 (k5_relat_1 X0 X1))) \quad (4)$$

Assume the following.

$$\forall X0.(v1_relat_1 (k4_relat_1 X0)) \wedge (v1_funct_1 (k4_relat_1 X0)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 X0) \Rightarrow (v1_relat_1 (k5_relat_1 \\ X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (k1_algspec1 \\ X0 X1 = k1_funct_4 (k6_partfun1 X0) (k5_relat_1 X1 X0)) \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (\\ & v1_funct_1 X1)) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow \\ & ((r1_tarski (k10_xtuple_0 X2) X0) \Rightarrow (k3_relat_1 X2 (k1_algspec1 \\ & X0 X1) = k3_relat_1 X2 (k1_funct_4 (k6_partfun1 X0) X1)))))) \end{aligned}$$