

t21_borsuk_7
(TMbcDf2Tt5BhyEtUT5629hjNNoa8ZSgdLab)

October 27, 2020

Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_borsuk_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. \forall X4. \forall X5. \\ & k9_xtuple_0 (k1_funct_4 (k1_funct_4 (k16_funcop_1 X0 X3) (k16_funcop_1 \\ & \quad X1 X4)) (k16_funcop_1 X2 X5)) = k1_enumset1 X0 X1 X2 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. k4_funct_4 X0 X1 X2 \\ & X3 = k1_funct_4 (k16_funcop_1 X0 X2) (k16_funcop_1 X1 X3) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. \forall X4. \forall X5. \\ & k1_borsuk_7 X0 X1 X2 X3 X4 X5 = k1_funct_4 (k4_funct_4 X0 X1 X3 X4) (\\ & \quad k16_funcop_1 X2 X5) \end{aligned} \tag{3}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. \forall X4. \forall X5. \\ & (v1_xcmplx_0 X5) \Rightarrow (k9_xtuple_0 (k1_borsuk_7 X0 X1 X5 X2 X3 X4) = k1_enumset1 \\ & \quad X0 X1 X5) \end{aligned}$$