

t21_bvfunc10

(TMWB5guj5kNtU2FCdcquvGQYuAwPy1vqYQs)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\
 & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
 & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\
 & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X3.((v1_funct_1 X3) \wedge \\
 & ((v1_funct_2 X3 X0 k6_margrel1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (((r1_bvfunc_1 X0 X1 X2) \wedge (\\
 & r1_bvfunc_1 X0 X2 X1)) \Rightarrow (r2_funct_2 X0 k6_margrel1 X1 X2)) \wedge (((r1_bvfunc_1 \\
 & X0 X1 X2) \wedge (r1_bvfunc_1 X0 X2 X3)) \Rightarrow (r1_bvfunc_1 X0 X1 X3)))))) \\
 & \tag{1}
 \end{aligned}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\
 & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
 & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\
 & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X3.((v1_funct_1 X3) \wedge \\
 & ((v1_funct_2 X3 X0 k6_margrel1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r1_bvfunc_1 X0 (k9_bvfunc_1 \\
 & X0 X1 X2) (k9_bvfunc_1 X0 X1 (k5_bvfunc_1 X0 X2 X3)))))) \\
 & \tag{2}
 \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge \\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r1_bvfunc_1 X0 (k2_bvfunc_1 \\ & X0 X1 X2) X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0) \wedge (((v1_funct_1 \\ & X1) \wedge ((v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \wedge ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k6_margrel1)))))) \Rightarrow ((v1_funct_1 (k9_bvfunc_1 X0 X1 X2)) \wedge \\ & (v1_funct_2 (k9_bvfunc_1 X0 X1 X2) X0 k6_margrel1) \wedge (m1_subset_1 \\ & (k9_bvfunc_1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0) \wedge (((v1_funct_1 \\ & X1) \wedge ((v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \wedge ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k6_margrel1)))))) \Rightarrow ((v1_funct_1 (k5_bvfunc_1 X0 X1 X2)) \wedge \\ & (v1_funct_2 (k5_bvfunc_1 X0 X1 X2) X0 k6_margrel1) \wedge (m1_subset_1 \\ & (k5_bvfunc_1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0) \wedge (((v1_funct_1 \\ & X1) \wedge ((v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \wedge ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k6_margrel1)))))) \Rightarrow ((v1_funct_1 (k2_bvfunc_1 X0 X1 X2)) \wedge \\ & (v1_funct_2 (k2_bvfunc_1 X0 X1 X2) X0 k6_margrel1) \wedge (m1_subset_1 \\ & (k2_bvfunc_1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0) \wedge (((v1_funct_1 \\ & X1) \wedge ((v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \wedge ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k6_margrel1)))))) \Rightarrow (k2_bvfunc_1 X0 X1 X2 = k2_bvfunc_1 X0 X2 \\ & X1)) \end{aligned} \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge \\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X3.((v1_funct_1 X3) \wedge \\ & ((v1_funct_2 X3 X0 k6_margrel1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r1_bvfunc_1 X0 (k2_bvfunc_1 \\ & X0 (k9_bvfunc_1 X0 X1 X2) (k9_bvfunc_1 X0 X2 X3)) (k9_bvfunc_1 X0 \\ & X2 (k5_bvfunc_1 X0 X3 X1)))) \end{aligned}$$