

t21_comseq_1

(TMPyc2DYBNk1PfJqPmVfT2SwAeTkrddbLdC)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k2_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k25_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_complex1 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k24_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ (\forall X1.(v1_xcmplx_0 X1) \Rightarrow (\forall X2.k1_funct_1 (k24_valued_1 \\ X0 X1) X2 = k3_xcmplx_0 X1 (k1_funct_1 X0 X2))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 np_1 X0 = X0) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\ X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow \\ (\forall X3.((v1_funct_1 X3) \wedge ((v1_funct_2 X3 X0 X1) \wedge (m1_subset_1 \\ X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow ((\forall X4.(X4 \in X0) \Rightarrow \\ (k1_funct_1 X2 X4 = k1_funct_1 X3 X4)) \Rightarrow (r2_relset_1 X0 X1 X2 X3))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.\forall X3.((v1_membered X1) \wedge \\ (((v1_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ X0 X1)))) \wedge (v1_xcmplx_0 X3))) \Rightarrow (k25_valued_1 X0 X1 X2 X3 = k24_valued_1 \\ X2 X3) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_valued_0 X0)))\Rightarrow(v1_xcmplx_0 (k1_funct_1 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(((\neg v1_xboole_0 X1)\wedge(v1_membered X1))\wedge(((v1_funct_1 X2)\wedge((v1_funct_2 X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))))\wedge(v1_xcmplx_0 X3)))\Rightarrow((v1_funct_1 (k24_valued_1 X2 X3))\wedge(v1_partfun1 (k24_valued_1 X2 X3) X0)) \quad (6)$$

Assume the following.

$$\neg v1_xboole_0 k2_numbers \quad (7)$$

Assume the following.

$$v1_membered k2_numbers \quad (8)$$

Assume the following.

$$m1_subset_1 k6_complex1 k2_numbers \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v1_membered X1)\wedge(((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\wedge(v1_xcmplx_0 X3)))\Rightarrow((v1_funct_1 (k25_valued_1 X0 X1 X2 X3))\wedge(m1_subset_1 (k25_valued_1 X0 X1 X2 X3) (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))) \quad (10)$$

Assume the following.

$$k6_complex1 = np_1 \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k2_numbers)\Rightarrow(v1_xcmplx_0 X0) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v1_partfun1 X2 X0)\Rightarrow(v1_funct_2 X2 X0 X1)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.(v1_membered X1)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_valued_0 X2)) \quad (15)$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 \ k5_numbers \ k2_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \ k5_numbers \ k2_numbers)))))) \Rightarrow \\ & (r2_relset_1 \ k5_numbers \ k2_numbers (k25_valued_1 \ k5_numbers \\ & \quad \quad \quad k2_numbers \ X0 \ k6_complex1) \ X0) \end{aligned}$$