

# t21\_connsp\_3 (TMJsqrNQaUPMotNNFy- BxJF4TVPqi4kCPvCC)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v2\_pre\_topc : \iota \Rightarrow o$  be given. Let  $l1\_pre\_topc : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $m1\_connsp\_3 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_connsp\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_tarski : \iota \Rightarrow \iota$  be given. Let  $v3\_connsp\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_setfam\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (r1\_tarski X0 (k3\_tarski X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X2))) \Rightarrow (m1\_subset\_1 X0 X2) \quad (2)$$

Assume the following.

$$\forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v2\_pre\_topc X0) \wedge (l1\_pre\_topc X0))) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (((v3\_connsp\_1 X1 X0) \wedge (X2 \in X1)) \Rightarrow (X1 = k1\_connsp\_1 X0 X2)))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k1\_zfmisc\_1 X0))) \Rightarrow (k5\_setfam\_1 X0 X1 = k3\_tarski X1) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k3\_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. (X2 \in X3) \wedge (X3 \in X0))) \quad (5)$$

Assume the following.

$$\forall X0. (l1\_pre\_topc X0) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow ((m1\_connsp\_3 X1 X0) \Leftrightarrow (\exists X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k1\_zfmisc\_1 (u1\_struct\_0 X0)))) \wedge (\forall X3. (m1\_subset\_1 X3 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow ((X3 \in X2) \Rightarrow (v3\_connsp\_1 X3 X0))) \wedge (X1 = k5\_setfam\_1 (u1\_struct\_0 X0) X2)))))) \quad (6)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v2\_pre\_topc X0) \wedge (l1\_pre\_topc \\ & X0))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 \\ & X0))) \Rightarrow (\forall X2.(m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (((X2 \in X1) \wedge \\ & (m1\_connsp\_3 X1 X0)) \Rightarrow (r1\_tarSKI (k1\_connsp\_1 X0 X2) X1)))) \end{aligned}$$