

t21_finseq_4
(TMGtgznpGXBYKVawd86qj28x6Gdf6GYLmf7)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k4_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow \\ & ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(X1 \in k10_xtuple_0 X0) \Rightarrow (k4_finseq_4 X0 X1 \in k4_finseq_1 \\ & X0)) \end{aligned} \quad (2)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (k4_finseq_1 X0 = k9_xtuple_0 X0) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1.((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (\\ & k1_relset_1 X0 X1 = k9_xtuple_0 X1) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 \\ & X0))) \Rightarrow (m1_subset_1 (k4_finseq_4 X0 X1) k5_numbers) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge \\ (v1_finseq_1 X0)))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ (\forall X1.(X1 \in k10_xtuple_0 X0) \Rightarrow ((r1_xxreal_0 np_1 (k4_finseq_4 \\ X0 X1)) \wedge (r1_xxreal_0 (k4_finseq_4 X0 X1) (k3_finseq_1 X0)))) \end{aligned}$$