

t21_goedelcp (TM- LafhX4G2ytLEQn5QDfQnsZMHfexSwVgnx)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $r1_henmodel : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r4_calcul_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \neg (X0 \in X1) \wedge (v1_xboole_0 X1) \quad (1)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((X1 = k9_finseq_1 X0) \Leftrightarrow ((k4_finseq_1 X1 = k2_finseq_1 np_1) \wedge (k10_xtuple_0 X1 = k1_tarski X0))) \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow ((k7_finseq_1 X0 \ k1_xboole_0 = X0) \wedge (k7_finseq_1 \ k1_xboole_0 X0 = X0)) \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 \ (k9_qc_lang1 \\ X0) \ (k3_cqc_lang X0)) \Rightarrow (\forall X2.(m2_finseq_1 X2 \ (k3_cqc_lang \\ X0)) \Rightarrow (r4_calcul_1 X0 \ (k8_finseq_1 \ (k3_cqc_lang X0) \ (k8_finseq_1 \\ (k3_cqc_lang X0) X2 \ (k12_finseq_1 \ (k3_cqc_lang X0) X1)) \ (k12_finseq_1 \\ (k3_cqc_lang X0) X1)))))) \quad (7) \end{aligned}$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (8)$$

Assume the following.

$$\forall X0.k9_finseq_1 X0 = k5_finseq_1 X0 \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_finseq_1 X1 X0) \wedge (m1_finseq_1 X2 X0)) \Rightarrow (k8_finseq_1 X0 X1 X2 = k7_finseq_1 X1 X2) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v5_relat_1 X1 X0)) \Rightarrow (k2_relset_1 X0 X1 = k10_xtuple_0 X1) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k12_finseq_1 X0 X1 = k5_finseq_1 X1) \quad (12)$$

Assume the following.

$$\forall X0.\exists X1.(m1_finseq_1 X1 X0) \wedge ((v1_relat_1 X1) \wedge ((v4_relat_1 X1 \ k5_numbers) \wedge ((v5_relat_1 X1 X0) \wedge ((v1_funct_1 X1) \wedge ((v1_xboole_0 X1) \wedge ((v1_finset_1 X1) \wedge (v1_finseq_1 X1))))))) \quad (13)$$

Assume the following.

$$\forall X0.v1_finseq_1 (k5_finseq_1 X0) \quad (14)$$

Assume the following.

$$\forall X0.(v1_relat_1 (k5_finseq_1 X0)) \wedge (v1_funct_1 (k5_finseq_1 X0)) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow (m2_finseq_1 (k12_finseq_1 X0 X1) X0) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow (X2 \in X1)) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_tarski X0)\Leftrightarrow(\forall X2.(X2 \in X1)\Leftrightarrow (X2 = X0)) \quad (18)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ (k3_cqc_lang X0)))\Rightarrow(\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0))\Rightarrow((r1_henmodel X0 X1 X2)\Leftrightarrow(\exists X3.(m2_finseq_1 \\ X3 (k3_cqc_lang X0))\wedge((r1_tarski (k2_relset_1 (k3_cqc_lang X0) \\ X3) X1)\wedge(r4_calcul_1 X0 (k8_finseq_1 (k3_cqc_lang X0) X3 (k12_finseq_1 \\ (k3_cqc_lang X0) X2)))))))))) \end{aligned} \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow(v5_relat_1 X1 X0) \quad (20)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ (k3_cqc_lang X0)))\Rightarrow(\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0))\Rightarrow((X2 \in X1)\Rightarrow(r1_henmodel X0 X1 X2)))) \end{aligned}$$