

## t21\_lfuzzy\_1

(TMUQvjnrhc2yXRanZHmf9u28JVh19xoAw)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_rcomp\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $v2\_lfuzzy\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_fuzzy\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_lfuzzy\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_fuzzy\_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_fuzzy\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned}
 & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. ((v5\_relat\_1 X1 (k1\_rcomp\_1 \\
 & k6\_numbers np\_1)) \wedge ((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 X0 k1\_numbers) \wedge \\
 & (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k1\_numbers))))) \Rightarrow \\
 & (\forall X2. ((v5\_relat\_1 X2 (k1\_rcomp\_1 k6\_numbers np\_1)) \wedge ( \\
 & (v1\_funct\_1 X2) \wedge ((v1\_funct\_2 X2 X0 k1\_numbers) \wedge (m1\_subset\_1 \\
 & X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k1\_numbers))))) \Rightarrow (\forall X3. \\
 & ((v5\_relat\_1 X3 (k1\_rcomp\_1 k6\_numbers np\_1)) \wedge ((v1\_funct\_1 \\
 & X3) \wedge ((v1\_funct\_2 X3 X0 k1\_numbers) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 \\
 & (k2\_zfmisc\_1 X0 k1\_numbers))))) \Rightarrow ((r1\_fuzzy\_1 X1 X2) \wedge (r1\_fuzzy\_1 \\
 & X1 X3)) \Rightarrow (r1\_fuzzy\_1 X1 (k1\_fuzzy\_1 X0 X2 X3))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
 & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. (\neg v1\_xboole\_0 X1) \Rightarrow \\
 & (\forall X2. ((v5\_relat\_1 X2 (k1\_rcomp\_1 k6\_numbers np\_1)) \wedge ( \\
 & (v1\_funct\_1 X2) \wedge ((v1\_funct\_2 X2 (k2\_zfmisc\_1 X0 X1) k1\_numbers) \wedge \\
 & (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1) \\
 & k1\_numbers))))) \Rightarrow (\forall X3. ((v5\_relat\_1 X3 (k1\_rcomp\_1 k6\_numbers \\
 & np\_1)) \wedge ((v1\_funct\_1 X3) \wedge ((v1\_funct\_2 X3 (k2\_zfmisc\_1 X0 X1) \\
 & k1\_numbers) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (k2\_zfmisc\_1 \\
 & X0 X1) k1\_numbers))))) \Rightarrow ((r1\_fuzzy\_1 X2 X3) \Rightarrow (r1\_fuzzy\_1 (k2\_fuzzy\_4 \\
 & X1 X0 X2) (k2\_fuzzy\_4 X1 X0 X3))))))
 \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\wedge(m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))))\Rightarrow((r2\_relset\_1 X0 X1 X2 X3)\Leftrightarrow(X2 = X3)) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0)\wedge(((v5\_relat\_1 X1 (k1\_rcomp\_1 k6\_numbers np\_1))\wedge((v1\_funct\_1 X1)\wedge((v1\_funct\_2 X1 X0 k1\_numbers)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k1\_numbers))))))\wedge((v5\_relat\_1 X2 (k1\_rcomp\_1 k6\_numbers np\_1))\wedge((v1\_funct\_1 X2)\wedge((v1\_funct\_2 X2 X0 k1\_numbers)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k1\_numbers))))))))\Rightarrow(k1\_lfuzzy\_1 X0 X1 X2 = k1\_fuzzy\_1 X0 X1 X2) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge(\neg v1\_xboole\_0 X1))\Rightarrow(\neg v1\_xboole\_0 (k2\_zfmisc\_1 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0)\wedge((\neg v1\_xboole\_0 X1)\wedge(((v5\_relat\_1 X2 (k1\_rcomp\_1 k6\_numbers np\_1))\wedge((v1\_funct\_1 X2)\wedge((v1\_funct\_2 X2 (k2\_zfmisc\_1 X1 X0) k1\_numbers)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (k2\_zfmisc\_1 X1 X0) k1\_numbers))))))\wedge(((v5\_relat\_1 (k2\_fuzzy\_4 X0 X1 X2) (k1\_rcomp\_1 k6\_numbers np\_1))\wedge((v1\_funct\_1 (k2\_fuzzy\_4 X0 X1 X2))\wedge((v1\_funct\_2 (k2\_fuzzy\_4 X0 X1 X2) (k2\_zfmisc\_1 X0 X1) k1\_numbers)\wedge(m1\_subset\_1 (k2\_fuzzy\_4 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1) k1\_numbers))))))))\Rightarrow \quad (6)$$

Assume the following.

$$\forall X0.(\neg v1\_xboole\_0 X0)\Rightarrow(\forall X1.(((v5\_relat\_1 X1 (k1\_rcomp\_1 k6\_numbers np\_1))\wedge((v1\_funct\_1 X1)\wedge((v1\_funct\_2 X1 (k2\_zfmisc\_1 X0 X0) k1\_numbers)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (k2\_zfmisc\_1 X0 X0) k1\_numbers))))))\Rightarrow((v2\_lfuzzy\_1 X1 X0)\Leftrightarrow(r2\_relset\_1 (k2\_zfmisc\_1 X0 X0) k1\_numbers (k2\_fuzzy\_4 X0 X0 X1) X1))) \quad (7)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v5\_relat\_1 X1 (k1\_rcomp\_1 \\ k6\_numbers np\_1)) \wedge ((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 (k2\_zfmisc\_1 \\ X0 X0) k1\_numbers) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ (k2\_zfmisc\_1 X0 X0) k1\_numbers)))))) \Rightarrow (\forall X2.((v5\_relat\_1 \\ X2 (k1\_rcomp\_1 k6\_numbers np\_1)) \wedge ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\ X2 (k2\_zfmisc\_1 X0 X0) k1\_numbers) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 (k2\_zfmisc\_1 X0 X0) k1\_numbers)))))) \Rightarrow (((v2\_lfuzzy\_1 \\ X2 X0) \wedge (r1\_fuzzy\_1 X2 X1)) \Rightarrow (r1\_fuzzy\_1 X2 (k1\_lfuzzy\_1 (k2\_zfmisc\_1 \\ X0 X0) X1 (k2\_fuzzy\_4 X0 X0 X1)))))) \end{aligned}$$