

t21_member_1 (TMKUYXpE- BrGSYi6t5tqXy1tW8cpAyB8VNUV)

October 27, 2020

Let $v2_membered : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k5_xxreal_3 : \iota \Rightarrow \iota$ be given. Let $k6_member_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Let $k2_member_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k7_numbers) \Rightarrow (k2_member_1 X0 = k5_xxreal_3 X0) \quad (2)$$

Assume the following.

$$\forall X0. (v2_membered X0) \Rightarrow (k6_member_1 X0 = ReplSep (toset (\lambda X1 : \iota. m1_subset_1 X1 k7_numbers)) (\lambda X1 : \iota. X1 \in X0) (\lambda X1 : \iota. k2_member_1 X1))) \quad (3)$$

Assume the following.

$$\forall X0. (v1_xxreal_0 X0) \Leftrightarrow (X0 \in k7_numbers) \quad (4)$$

Theorem 1

$$\forall X0. (v2_membered X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow ((X1 \in X0) \Rightarrow (k5_xxreal_3 X1 \in k6_member_1 X0)))$$