

t21_moebius1

(TMcD2Au8rZzxr5AHTTHdsQiKmRkpVQBiBoo)

October 27, 2020

Let $v1_moebius1 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_nat_d : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $v1_int_2 : \iota \Rightarrow o$ be given. Let $np_2 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k1_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((r1_nat_d X0 k6_numbers) \wedge (r1_nat_d np_1 X0)) \quad (1)$$

Assume the following.

$$v1_int_2 np_2 \quad (2)$$

Assume the following.

$$m1_subset_1 k1_xboole_0 k4_ordinal1 \quad (3)$$

Assume the following.

$$((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \quad (4)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (v7_ordinal1 X1)) \Rightarrow (v7_ordinal1 (k1_newton X0 X1)) \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow((v1_moebius1\ X0)\Leftrightarrow(\exists X1.(v7_ordinal1\ X1)\wedge(v1_int_2\ X1)\wedge(r1_nat_d\ (k1_newton\ X1\ np_2)\ X0))) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k4_ordinal1)\Rightarrow(v7_ordinal1\ X0) \quad (9)$$

Theorem 1 *v1_moebius1 k6_numbers.*