

t21_pdiff_3 (TM- Rzb9vNsaDCvEkPoDQnuzWLrfMsURDTqUv)

October 27, 2020

Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_pdiff_3 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r3_pdiff_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k3_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_pdiff_3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k11_pdiff_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_pdiff_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m2_finseq_2 X0 k1_numbers (k1_euclid np_2)) \Rightarrow (\forall X1. \\ & ((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 (\\ & k1_euclid np_2) k1_numbers)))) \Rightarrow ((r1_pdiff_3 X1 X0) \Leftrightarrow (r3_pdiff_1 \\ & np_2 np_1 (k1_pdiff_3 np_1 np_2 X1) X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v1_xboole_0 X0) \wedge (m2_subset_1 X0 k1_numbers k5_numbers)) \Rightarrow \\ & (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (\forall X2. \\ & ((v1_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 (\\ & k1_euclid X0) k1_numbers)))) \Rightarrow (\forall X3.((v1_funct_1 X3) \wedge (\\ & m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 (k1_euclid X0) k1_numbers)))) \Rightarrow \\ & (\forall X4.(m2_finseq_2 X4 k1_numbers (k1_euclid X0)) \Rightarrow (((r3_pdiff_1 \\ & X0 X1 X2 X4) \wedge (r3_pdiff_1 X0 X1 X3 X4)) \Rightarrow ((r3_pdiff_1 X0 X1 (k3_valued_1 \\ & (k1_euclid X0) k1_numbers k1_numbers X2 X3) X4) \wedge (k11_pdiff_1 X0 \\ & X1 (k3_valued_1 (k1_euclid X0) k1_numbers k1_numbers X2 X3) X4 = \\ & k7_real_1 (k11_pdiff_1 X0 X1 X2 X4) (k11_pdiff_1 X0 X1 X3 X4)))))))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m2_finseq_2 X0 k1_numbers (k1_euclid np_2)) \Rightarrow (\forall X1. \\ & ((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 (\\ & k1_euclid np_2) k1_numbers)))) \Rightarrow ((r1_pdiff_3 X1 X0) \Rightarrow (k2_pdiff_3 \\ & X1 X0 = k11_pdiff_1 np_2 np_1 (k1_pdiff_3 np_1 np_2 X1) X0))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$\neg v1_xboole_0 np_2 \quad (5)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((m1_subset_1 X0 k5_numbers) \wedge \\ & (((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 k5_numbers)) \wedge ((v1_funct_1 \\ & X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 (k1_euclid X1) \\ & k1_numbers)))))) \Rightarrow ((v1_funct_1 (k1_pdiff_3 X0 X1 X2)) \wedge ((v1_funct_2 \\ & (k1_pdiff_3 X0 X1 X2) (k1_euclid X1) k1_numbers) \wedge (m1_subset_1 \\ & (k1_pdiff_3 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 (k1_euclid X1) \\ & k1_numbers)))))) \end{aligned} \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m2_finseq_2 X0 k1_numbers (k1_euclid np_2)) \Rightarrow (\forall X1. \\ & ((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 (\\ & k1_euclid np_2) k1_numbers)))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 (k1_euclid np_2) k1_numbers)))) \Rightarrow \\ & (((r1_pdiff_3 X1 X0) \wedge (r1_pdiff_3 X2 X0)) \Rightarrow ((r3_pdiff_1 np_2 np_1 \\ & (k3_valued_1 (k1_euclid np_2) k1_numbers k1_numbers (k1_pdiff_3 \\ & np_1 np_2 X1) (k1_pdiff_3 np_1 np_2 X2)) X0) \wedge (k11_pdiff_1 np_2 \\ & np_1 (k3_valued_1 (k1_euclid np_2) k1_numbers k1_numbers (k1_pdiff_3 \\ & np_1 np_2 X1) (k1_pdiff_3 np_1 np_2 X2)) X0 = k7_real_1 (k2_pdiff_3 \\ & X1 X0) (k2_pdiff_3 X2 X0)))))) \end{aligned}$$