

t21_pre_circ
(TMFw4Q8LBqdGceQoDjohUahWkoWDZnZz6zD)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r2_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $r2_wellord2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (k1_card_1 X0 = k1_card_1 (k9_xtuple_0 X0)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r2_wellord2 X0 X1) \Leftrightarrow (k1_card_1 X0 = k1_card_1 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r2_wellord2 X0 X1) \Leftrightarrow (r2_tarski X0 X1) \quad (3)$$

Theorem 1

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r2_tarski (k9_xtuple_0 X0) (k9_xtuple_0 X1)) \Leftrightarrow (r2_tarski X0 X1)))$$