

t21_rat_1

(TMRx57QvUf4F1JPXo5z2K7P2gu9pBokWdNM)

October 27, 2020

Let $v1_rat_1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_rat_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_int_1 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k2_rat_1 : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X0)) \Rightarrow (X0 = X1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_rat_1 X0) \Rightarrow (((k2_rat_1 X0 = X0) \vee (k1_rat_1 X0 = np_1)) \Rightarrow (v1_int_1 X0)) \quad (2)$$

Assume the following.

$$\forall X0.(v1_rat_1 X0) \Rightarrow ((v1_int_1 X0) \Rightarrow ((k1_rat_1 X0 = np_1) \wedge (k2_rat_1 X0 = X0))) \quad (3)$$

Assume the following.

$$\forall X0.(v1_rat_1 X0) \Rightarrow (r1_xxreal_0 np_1 (k1_rat_1 X0)) \quad (4)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (5)$$

Assume the following.

$$r1_xxreal_0 np_1 np_1 \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (8)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (9)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ X2 X0 X1) \Rightarrow (m1_subset_1 X2 X0)) \end{aligned} \quad (10)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (11)$$

Assume the following.

$$\forall X0. (v1_rat_1 X0) \Rightarrow (m2_subset_1 (k1_rat_1 X0) k1_numbers \\ k5_numbers) \quad (12)$$

Assume the following.

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (13)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (14)$$

Theorem 1

$$\begin{aligned} \forall X0. (v1_rat_1 X0) \Rightarrow ((\neg(\neg r1_xxreal_0 (k1_rat_1 X0) np_1) \wedge \\ (v1_int_1 X0)) \wedge (\neg(\neg v1_int_1 X0) \wedge (r1_xxreal_0 (k1_rat_1 X0) np_1))) \end{aligned}$$