

# t21\_rlvect\_1 (TMQvr- cXyZQFP15mtjBRF9BJuYB6mR5WV1wx)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $l2\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k5\_algstr\_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k4\_algstr\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_algstr\_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v3\_rlvect\_1 \\ X0) \wedge ((v4\_rlvect\_1 X0) \wedge (l2\_algstr\_0 X0)))))) \Rightarrow (\forall X1.(m1\_subset\_1 \\ X1 (u1\_struct\_0 X0)) \Rightarrow (k4\_algstr\_0 X0 (k4\_algstr\_0 X0 X1) = X1)) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((l2\_algstr\_0 X0) \wedge (m1\_subset\_1 X1 (u1\_struct\_0 \\ X0))) \Rightarrow (m1\_subset\_1 (k4\_algstr\_0 X0 X1) (u1\_struct\_0 X0)) \quad (2)$$

Assume the following.

$$\forall X0. (l2\_algstr\_0 X0) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (u1\_struct\_0 \\ X0)) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (k5\_algstr\_0 \\ X0 X1 X2 = k1\_algstr\_0 X0 X1 (k4\_algstr\_0 X0 X2)))) \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0. ((\neg v2\_struct\_0 X0) \wedge (l2\_algstr\_0 X0)) \Rightarrow (\forall X1. \\ (m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 \\ X0) \wedge (v13\_algstr\_0 X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (u1\_struct\_0 \\ X0)) \Rightarrow ((X2 = k4\_algstr\_0 X0 X1) \Leftrightarrow (k1\_algstr\_0 X0 X1 X2 = k4\_struct\_0 \\ X0)))))) \end{aligned} \quad (4)$$

## Theorem 1

$$\begin{aligned} \forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v3\_rlvect\_1 \\ X0) \wedge ((v4\_rlvect\_1 X0) \wedge (l2\_algstr\_0 X0)))))) \Rightarrow (\forall X1. (m1\_subset\_1 \\ X1 (u1\_struct\_0 X0)) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (u1\_struct\_0 \\ X0)) \Rightarrow ((k5\_algstr\_0 X0 X1 X2 = k4\_struct\_0 X0) \Rightarrow (X1 = X2)))) \end{aligned}$$