

t21\_xxreal\_2  
(TMU1fSfLa8frJAUXfg2VjYk46er8ibMAEqZ)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m1\_xxreal\_2 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v2\_membered : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow ((X0 \in k1\_xxreal\_1 X1 X2) \Leftrightarrow ((r1\_xxreal\_0 X1 X0) \wedge \\ & (r1\_xxreal\_0 X0 X2)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow (v2\_membered (k1\_xxreal\_1 X0 X1)) \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v2\_membered X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (( \\ & m1\_xxreal\_2 X1 X0) \Leftrightarrow (\forall X2.(v1\_xxreal\_0 X2) \Rightarrow ((X2 \in X0) \Rightarrow (r1\_xxreal\_0 \\ & X2 X1)))))) \end{aligned} \tag{3}$$

**Theorem 1**

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (m1\_xxreal\_2 X0 (k1\_xxreal\_1 X1 X0)))$$