

t223_xxreal_1

(TMMH1nxE4btKxEJhvmF7DvKZJxjtufCD7Tc)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((X0 \in k4_xxreal_1 X1 X2) \Leftrightarrow ((\neg r1_xxreal_0 X0 X1) \wedge \\ & (\neg r1_xxreal_0 X2 X0)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(X2 = k2_xboole_0 X0 X1) \Leftrightarrow (\forall X3. \\ & (X3 \in X2) \Leftrightarrow ((X3 \in X0) \vee (X3 \in X1))) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow (\neg(r1_xxreal_0 \\ & X0 X1) \wedge ((r1_xxreal_0 X1 X2) \wedge (X0 \in k2_xboole_0 (k4_xxreal_1 X1 X2) \\ & (k4_xxreal_1 X2 X3)))))))))) \end{aligned}$$