

t226_xxreal_1 (TM-
cdHv5FvaKeBK8rWmk41LkqpdtkTAC4SdG)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (((X0 \in k1_numbers) \wedge (r1_xxreal_0 X0 X1)) \Rightarrow ((\\ & r1_xxreal_0 X2 X1) \vee (X1 \in k1_numbers)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((X0 \in k2_xxreal_1 X1 X2) \Leftrightarrow ((r1_xxreal_0 X1 X0) \wedge \\ & (\neg r1_xxreal_0 X2 X0)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (v2_membered (k2_xxreal_1 X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0.(v2_membered X0) \Rightarrow (\forall X1.(r1_tarski X0 X1) \Leftrightarrow (\forall X2. (v1_xxreal_0 X2) \Rightarrow ((X2 \in X0) \Rightarrow (X2 \in X1)))) \quad (4)$$

Theorem 1

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((X0 \in k1_numbers) \Rightarrow (r1_tarski (k2_xxreal_1 X0 X1) k1_numbers)))$$