

t229_xreal_1

(TMP5baqD91wdXRNtQhkWh4eEiG93fHxb8uF)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\neg \\ & (\neg r1_xxreal_0 X1 X0) \wedge (\forall X2.(v1_xxreal_0 X2) \Rightarrow (\neg (\neg r1_xxreal_0 \\ & X2 X0) \wedge (\neg r1_xxreal_0 X1 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (\\ & (r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0)) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((\forall X3.(v1_xxreal_0 X3) \Rightarrow (\neg (\neg r1_xxreal_0 \\ & X3 X0) \wedge (\neg r1_xxreal_0 X1 X3) \wedge (\neg r1_xxreal_0 X3 X2)))) \Rightarrow ((r1_xxreal_0 \\ & X1 X0) \vee (r1_xxreal_0 X1 X2)))))) \end{aligned}$$