

## t22\_bvfunc\_4

(TMHMX3hCQzhTSUM2dRxbLGHS7BSQVbQhLeT)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_margrel1 : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_bvfunc\_2 : \iota \Rightarrow \iota$  be given. Let  $m1\_eqrel\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r3\_bvfunc\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k7\_bvfunc\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r2\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_partit1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge \\ (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2.(m1\_subset\_1 X2 \\ (k1\_zfmisc\_1 (k1\_bvfunc\_2 X0))) \Rightarrow (\forall X3.(m1\_eqrel\_1 X3 X0) \Rightarrow \\ ((r3\_bvfunc\_2 X0 X1 X2 X3) \Rightarrow (r1\_bvfunc\_1 X0 (k7\_bvfunc\_2 X0 X1 X2 \\ X3) X1)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge \\ (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge \\ ((v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X3.((v1\_funct\_1 X3) \wedge \\ ((v1\_funct\_2 X3 X0 k6\_margrel1) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (((r1\_bvfunc\_1 X0 X1 X2) \wedge \\ (r1\_bvfunc\_1 X0 X2 X1)) \Rightarrow (r2\_funct\_2 X0 k6\_margrel1 X1 X2) \wedge ((r1\_bvfunc\_1 \\ X0 X1 X2) \wedge (r1\_bvfunc\_1 X0 X2 X3)) \Rightarrow (r1\_bvfunc\_1 X0 X1 X3)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 \\ (k1\_partit1 X0))) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\ X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ X0 k6\_margrel1)))))) \Rightarrow (\forall X3.(m1\_eqrel\_1 X3 X0) \Rightarrow (r1\_bvfunc\_1 \\ X0 X2 (k7\_bvfunc\_2 X0 X2 X1 X3)))))) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.k1\_bvfunc\_2 X0 = k1\_partit1 X0 \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1\_xboole\_0 X0) \wedge \\ & (((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 \\ & X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \wedge ((m1\_subset\_1 \\ & X2 (k1\_zfmisc\_1 (k1\_bvfunc\_2 X0))) \wedge (m1\_eqrel\_1 X3 X0)))) \Rightarrow ((v1\_funct\_1 \\ & (k7\_bvfunc\_2 X0 X1 X2 X3)) \wedge ((v1\_funct\_2 (k7\_bvfunc\_2 X0 X1 X2 X3) \\ & X0 k6\_margrel1) \wedge (m1\_subset\_1 (k7\_bvfunc\_2 X0 X1 X2 X3) (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \end{aligned} \quad (5)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge ( \\ & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2.(m1\_subset\_1 X2 \\ & (k1\_zfmisc\_1 (k1\_bvfunc\_2 X0))) \Rightarrow (\forall X3.(m1\_eqrel\_1 X3 X0) \Rightarrow \\ & (\forall X4.(m1\_eqrel\_1 X4 X0) \Rightarrow ((r3\_bvfunc\_2 X0 X1 X2 X3) \Rightarrow (r1\_bvfunc\_1 \\ & X0 (k7\_bvfunc\_2 X0 X1 X2 X3) (k7\_bvfunc\_2 X0 X1 X2 X4)))))) \end{aligned}$$