

t22_fib_num2 (TMXtqRgV- zoL1Fg8tMWDHZ3qF7pK2fCa9WfF)

October 27, 2020

Let $k1_pre_ff : \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $np_2 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Assume the following.

$$\begin{aligned} & (k1_pre_ff\ k6_numbers = k6_numbers) \wedge ((k1_pre_ff\ np_1 = np_1) \wedge \\ & (\forall X0.(v7_ordinal1\ X0) \Rightarrow (k1_pre_ff\ (k2_nat_1\ (k1_nat_1 \\ & X0\ np_1)\ np_1) = k2_nat_1\ (k1_pre_ff\ X0)\ (k1_pre_ff\ (k1_nat_1 \\ & X0\ np_1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0\ np_2) \wedge (m2_subset_1\ np_2\ k1_numbers\ k5_numbers)) \wedge \\ & ((m1_subset_1\ np_2\ k5_numbers) \wedge (m1_subset_1\ np_2\ k1_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0\ np_1) \wedge (m2_subset_1\ np_1\ k1_numbers\ k5_numbers)) \wedge \\ & ((m1_subset_1\ np_1\ k5_numbers) \wedge (m1_subset_1\ np_1\ k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$k2_xcmplx_0\ np_2\ np_1 = np_3 \quad (4)$$

Assume the following.

$$k2_xcmplx_0\ np_1\ np_1 = np_2 \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(m1_subset_1 X1 k5_numbers))\Rightarrow(k1_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (8)$$

Assume the following.

$$k1_pre_ff np_2 = np_1 \quad (9)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (10)$$

Theorem 1 $k1_pre_ff np_3 = np_2$.