

t22\_finseqop  
(TMJpdr46WSGuoUqpGygpSQfujRvtNnJomdz)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $v3\_card\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_finseqop : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_funcop\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_partfun1 : \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k4\_relat\_1 : \iota \Rightarrow \iota$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. (v1\_relat\_1 X1) \Rightarrow ((r1\_tarski (k10\_xtuple\_0 X1) X0) \Rightarrow (k3\_relat\_1 X1 (k4\_relat\_1 X0) = X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (\forall X2. ((v1\_relat\_1 X2) \wedge (v1\_funct\_1 X2)) \Rightarrow (\forall X3. k3\_relat\_1 X1 (k5\_funcop\_1 X2 X3 X0) = k5\_funcop\_1 X2 X3 (k3\_relat\_1 X1 X0)))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (4)$$

Assume the following.

$$\forall X0. k6\_partfun1 X0 = k4\_relat\_1 X0 \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.\forall X4.\forall X5. \\ & ((\neg v1\_xboole\_0 X0)\wedge((\neg v1\_xboole\_0 X1)\wedge((\neg v1\_xboole\_0 X2)\wedge( \\ & ((v1\_funct\_1 X3)\wedge((v1\_funct\_2 X3 (k2\_zfmisc\_1 X0 X1) X2)\wedge(m1\_subset\_1 \\ & X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1) X2))))))\wedge((m1\_subset\_1 \\ & X4 X0)\wedge(m1\_finseq\_1 X5 X1))))))\Rightarrow(k3\_finseqop X0 X1 X2 X3 X4 X5 = k5\_funcop\_1 \\ & X3 X4 X5) \end{aligned} \tag{6}$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge(v5\_relat\_1 X1 X0))\Rightarrow(k2\_relset\_1 X0 X1 = k10\_xtuple\_0 X1) \tag{7}$$

Assume the following.

$$\forall X0.\forall X1.v1\_relat\_1 (k2\_zfmisc\_1 X0 X1) \tag{8}$$

Assume the following.

$$\forall X0.(v1\_relat\_1 (k4\_relat\_1 X0))\wedge(v1\_funct\_1 (k4\_relat\_1 X0)) \tag{9}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(m2\_finseq\_1 X1 X0)\Rightarrow((v1\_funct\_1 X1)\wedge( \\ & (v1\_finseq\_1 X1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers \\ & X0)))))) \end{aligned} \tag{10}$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0)\Rightarrow((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1))) \tag{11}$$

Assume the following.

$$\forall X0.v1\_relat\_1 (k4\_relat\_1 X0) \tag{12}$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge(v5\_relat\_1 X1 X0))\Rightarrow(m1\_subset\_1 (k2\_relset\_1 X0 X1) (k1\_zfmisc\_1 X0)) \tag{13}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \tag{14}$$

Assume the following.

$$\forall X0.(v1\_relat\_1 X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0))\Rightarrow(v1\_relat\_1 X1)) \tag{15}$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.(\neg v1\_xboole\_0 X1) \Rightarrow \\ & (\forall X2.(\neg v1\_xboole\_0 X2) \Rightarrow (\forall X3.(m1\_subset\_1 X3 X0) \Rightarrow \\ & (\forall X4.(v7\_ordinal1 X4) \Rightarrow (\forall X5.((v1\_funct\_1 X5) \wedge ( \\ & (v1\_funct\_2 X5 (k2\_zfmisc\_1 X0 X2) X1) \wedge (m1\_subset\_1 X5 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 (k2\_zfmisc\_1 X0 X2) X1)))))) \Rightarrow (\forall X6.((v3\_card\_1 \\ & X6 X4) \wedge (m2\_finseq\_1 X6 X2)) \Rightarrow (k3\_finseqop X0 X2 X1 X5 X3 X6 = k3\_relat\_1 \\ & X6 (k5\_funcop\_1 X5 X3 (k6\_partfun1 X2)))))))))) \end{aligned}$$