

t22\_kurato\_0  
(TMczc23crnbZGPo6jDcK7AriMR5tDZC95fb)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k9\_setfam\_1 : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k3\_funct\_1 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X2))) \Rightarrow (m1\_subset\_1 X0 X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. k9\_setfam\_1 X0 = k1\_zfmisc\_1 X0 \quad (3)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. (X1 = k10\_xtuple\_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. (X3 \in k9\_xtuple\_0 X0) \wedge (X2 = k1\_funct\_1 X0 X3)))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (v1\_relat\_1 X1) \Rightarrow ((v5\_relat\_1 X1 X0) \Leftrightarrow (r1\_tarski (k10\_xtuple\_0 X1) X0)) \quad (5)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow ((v3\_funct\_1 X0) \Rightarrow ((v1\_xboole\_0 X0) \vee (\forall X1. (X1 = k3\_funct\_1 X0) \Leftrightarrow (\exists X2. (X2 \in k9\_xtuple\_0 X0) \wedge (X1 = k1\_funct\_1 X0 X2)))))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ k5\_numbers (k9\_setfam\_1 X0))))\Rightarrow(((v1\_funct\_1 X1)\wedge(v1\_funct\_2 \\ X1 k5\_numbers (k9\_setfam\_1 X0)))\Rightarrow((\neg v1\_xboole\_0 X1)\wedge((v1\_funct\_1 \\ X1)\wedge(v1\_funct\_2 X1 k5\_numbers (k9\_setfam\_1 X0)))))) \end{aligned} \quad (9)$$

**Theorem 1**

$$\begin{aligned} \forall X0.\forall X1.((v1\_funct\_1 X1)\wedge((v1\_funct\_2 X1 k5\_numbers \\ (k9\_setfam\_1 X0))\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ k5\_numbers (k9\_setfam\_1 X0))))))\Rightarrow((v3\_funct\_1 X1)\Rightarrow(m1\_subset\_1 \\ (k3\_funct\_1 X1) (k1\_zfmisc\_1 X0))) \end{aligned}$$