

t22_nfcont_1

(TMcQFco2diHYcTs6UztFPT4rcBngMwFq96z)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_normsp_0 : \iota \Rightarrow o$ be given. Let $v4_normsp_0 : \iota \Rightarrow o$ be given. Let $v2_normsp_1 : \iota \Rightarrow o$ be given. Let $l1_normsp_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r4_nfcont_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_nfcont_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k5_relat_1 (k5_relat_1 X1 X0) X0 = k5_relat_1 X1 X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k9_xtuple_0 (k5_relat_1 X1 X0) = k3_xboole_0 (k9_xtuple_0 X1) X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Rightarrow (k3_xboole_0 X0 X1 = X0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski (k3_xboole_0 X0 X1) X0 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. ((v1_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \Rightarrow (k2_partfun1 X0 X1 X2 X3 = k5_relat_1 X2 X3) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((v1_funct_1 (k2_partfun1 X0 X1 X2 X3))\wedge(m1_subset_1 (k2_partfun1 X0 X1 X2 X3) (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0)\wedge((v13_algstr_0 X0)\wedge((v2_rlvect_1 X0)\wedge((v3_rlvect_1 X0)\wedge((v4_rlvect_1 X0)\wedge((v5_rlvect_1 X0)\wedge \\ & ((v6_rlvect_1 X0)\wedge((v7_rlvect_1 X0)\wedge((v8_rlvect_1 X0)\wedge((v3_normsp_0 X0)\wedge((v4_normsp_0 X0)\wedge((v2_normsp_1 X0)\wedge(l1_normsp_1 X0))))))))))\Rightarrow \\ & (\forall X1.((v1_funct_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 (u1_struct_0 X0) k1_numbers))))\Rightarrow(\forall X2.(r4_nfcont_1 X0 X1 X2)\Leftrightarrow((r1_tarSKI X2 (k1_relset_1 (u1_struct_0 X0) X1))\wedge(\forall X3. \\ & (m1_subset_1 X3 (u1_struct_0 X0))\Rightarrow((X3 \in X2)\Rightarrow(r2_nfcont_1 X0 (k2_partfun1 (u1_struct_0 X0) k1_numbers X1 X2) X3)))))) \quad (8) \end{aligned}$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (11)$$

Theorem 1

$$\begin{aligned} & \forall X0.\forall X1.((\neg v2_struct_0 X1)\wedge((v13_algstr_0 X1)\wedge \\ & ((v2_rlvect_1 X1)\wedge((v3_rlvect_1 X1)\wedge((v4_rlvect_1 X1)\wedge((v5_rlvect_1 X1)\wedge \\ & X1)\wedge((v6_rlvect_1 X1)\wedge((v7_rlvect_1 X1)\wedge((v8_rlvect_1 X1)\wedge \\ & ((v3_normsp_0 X1)\wedge((v4_normsp_0 X1)\wedge((v2_normsp_1 X1)\wedge(l1_normsp_1 X1))))))))))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge(m1_subset_1 X2 \\ & (k1_zfmisc_1 (k2_zfmisc_1 (u1_struct_0 X1) k1_numbers))))\Rightarrow \\ & ((r4_nfcont_1 X1 X2 X0)\Leftrightarrow(r4_nfcont_1 X1 (k2_partfun1 (u1_struct_0 X1) k1_numbers X2 X0) X0))) \end{aligned}$$